

MAPLE SUBROUTINES FOR COMPUTING MILNOR AND TYURINA NUMBERS OF HYPERSURFACE SINGULARITIES WITH APPLICATION TO ARNOL'D ADJACENCIES.

MICHELE ROSSI AND LEA TERRACINI

ABSTRACT. In the present paper MAPLE subroutines computing Milnor and Tyurina numbers of an isolated algebraic hypersurface singularity are presented and described in detail. They represent examples, and perhaps the first ones, of a MAPLE implementation of *local monomial ordering*.

As an application, the last section is devoted to writing down equations of algebraic stratifications of Kuranishi spaces of simple Arnol'd singularities: they geometrically represent, by means of inclusions of algebraic subsets, the partial ordering on classes of simple singularities induced by the *adjacency* relation.

CONTENTS

1. Milnor number of an isolated hypersurface singularity	3
1.1. Milnor number from the algebraic point of view	4
2. Tyurina number of an isolated hypersurface singularity	4
2.1. Deformations of complex spaces	4
2.2. Deformations of an i.h.s.	5
3. Milnor and Tyurina numbers of a polynomial	5
3.1. Milnor and Tyurina numbers of weighted homogeneous polynomials	6
3.2. An example: weighted homogeneous cDV singularities	8
3.3. The algebraic computation via Gröbner basis	9
4. Monomial Ordering	9
4.1. Localizations in $\mathbb{C}[\mathbf{x}]$ and rings implemented by monomial orders	10
5. MAPLE subroutines in detail	11
5.1. The subroutines	12
5.2. Some user friendly examples	17
5.3. Optional input: some more subtle utilities	20
5.4. An efficiency remark: global to local subroutines	23
6. Application: adjacencies of Arnol'd simple singularities	27
6.1. Outline of the following results	30
6.2. Simple singularities of A_n type.	31
6.3. Simple singularities of D_n type.	35
6.4. Simple singularities of E_6 type.	42
6.5. Simple singularities of E_7 type.	48

This work has been developed despite the effects of the Italian law 133/08 (<http://groups.google.it/group/scienceaction>). This law drastically reduces public funds to public Italian universities, which is particularly dangerous for scientific free research, and it will prevent young researchers from getting a position, either temporary or tenured, in Italy. The authors are protesting against this law to obtain its repeal.

6.6. Simple singularities of E_8 type.	57
6.7. A list of very special adjacencies	66
References	68

Two basic invariants of a complex analytic complete intersection singularity $p \in \overline{U} = \mathbf{f}^{-1}(0)$ are its *Milnor number* $\mu(p)$ and its *Tyurina number* $\tau(p)$. The former essentially "counts the number" of vanishing cycles in the intermediate cohomology of a nearby smoothing $U_t = \mathbf{f}^{-1}(t)$ of $\overline{U}$, which actually turns out to be the *multiplicity of p as a critical point* of the map $\mathbf{f}$. The latter "counts the dimension" of the base space of a versal deformation of $p \in \overline{U}$, which actually turn out to be the *multiplicity of p as a singularity* of the complex space $\overline{U}$. After the Looijenga–Steenbrink Theorem [17] it is a well known fact that $\tau(p) \leq \mu(p)$.

The purpose of the present paper is to present MAPLE subroutines [1] allowing to compute those invariants in the case of an isolated algebraic hypersurface singularity (i.h.s.). In fact, from the computational point of view, their calculation could be very intricate to end up and a computer employment may be needed in most situations. Let us underline that the actual originality of our procedure is not so much based on the effective computation of those invariants as on its implementation with a very interdisciplinary and worldwide diffused and known math software like MAPLE is. In fact computer algebra packages computing those singularities' invariants already exists (one for all is SINGULAR [2]). But our hope is that routines here presented could be useful to all those who are interesting, for any reason, in a concrete evaluation of those invariants without being so much motivated for learning an entire computer algebra package. Once implemented, the present routine is so easy that even an undergraduate student may use it!

In other words we believe that the present routine could be an interesting and, as far as we know, the first example of a MAPLE implementation of *local monomial ordering* (l.m.o.). In fact term orders, i.e. usual monomial ordering (recently called *global* in contrast with the word *local*), are already implemented in MAPLE with the command `MonomialOrder` in the `Groebner` package. Actually user-defined term orders are also allowed and, in particular, l.m.o.'s can be easily defined as the *opposite* of a standard g.m.o. (pure lexicographic, graded lexicographic, reverse ... etc.). The problem is that Buchberger S-procedure may not end up determining a *normal form* since a l.m.o. is not a well-order on the contrary of g.m.o.'s. Then the present MAPLE routines start with an implementation of the Mora algorithm for determining a *weak normal form* and then monomial bases of Milnor and Tyurina ideals in the complex ring of convergent power series $\mathbb{C}\{x_1, \dots, x_{n+1}\}$ (see [20], [13] Algorithm 1.7.6, [8] Algorithm 9.22). As a consequence a monomial basis of the Kuranishi space, parameterizing small versal deformations of the given i.h.s., is obtained, allowing to concretely write down these deformations even for more intricate cases. Actually we get procedures which are able to perform calculation in $(\mathbb{C}[\lambda])[\mathbf{x}]$, where $\lambda = (\lambda_1, \dots, \lambda_r)$ is an r -tuple of parameters e.g. coordinates of the Kuranishi space.

An interesting application of this latter feature is that of *writing down equations* of algebraic stratifications in Kuranishi spaces of Arnol'd simple singularities,

giving an *explicit* geometric interpretation, by means of inclusions of algebraic subsets, of *Arnol'd's adjacency partial order relation* over classes of simple singularities (see Section 6). This section ends up with an explicit list of the most specialized 1-parameter deformations of simple singularities realizing adjacencies between distinct classes of simple singularities (see 6.7).

Acknowledgements. We are greatly indebt with G. M. Greuel who timely pointed out to us serious mistakes in the first version of the this note. His concise and sharp remarks considerably helped us to improve our routine and the final product. Authors would also like to thank A. Albano for enlightening conversations.

1. MILNOR NUMBER OF AN ISOLATED HYPERSURFACE SINGULARITY

From the topological point of view a *good representative* $\overline{U}$ of an *isolated hypersurface singularity (i.h.s.)* is the zero locus of a holomorphic map

$$(1) \quad f : \mathbb{C}^{n+1} \longrightarrow \mathbb{C} \quad , \quad n \geq 0$$

admitting an isolated critical point in $0 \in \mathbb{C}^4$.

Set $U_T := f^{-1}(T)$ where T is a small enough neighborhood of $0 \in \mathbb{C}$. Then we can assume that f is a submersion over $U_T \setminus \{0\}$: therefore $\overline{U} = f^{-1}(0) = U_0$ and U_t is a *local smoothing* of $\overline{U}$, for $0 \neq t \in T$.

Definition 1.1. Let $X \subset \mathbb{C}^m$ be a subset. The following subset of $\mathbb{C}^m$

$$\text{Cn}_0(X) := \{tx \mid \forall t \in [0, 1] \subset \mathbb{R} \, , \, \forall x \in X\}$$

will be called the *cone projecting* X .

Theorem 1.2 (Local topology of a isolated hypersurface singularity, [18] Theorem 2.10, Theorem 5.2). *Let D_ε denote the closed ball of radius $\varepsilon > 0$, centered in $0 \in \mathbb{C}^{n+1}$, whose boundary is the $2n+1$ -dimensional sphere S_ε^{2n+1} . Then, for ε small enough, the intersection*

$$\overline{B}_\varepsilon := \overline{U} \cap D_\varepsilon$$

is homeomorphic to the cone $\text{Cn}_0(K)$ projecting $K := \overline{U} \cap S_\varepsilon^{2n+1}$, which is called the knot or link of the singularity $0 \in \overline{U}$.

Theorem–Definition 1.3 (Local homology type of the smoothing [18], Theorems 5.11, 6.5, 7.2). *Set $\tilde{U} := U_t$ for some $0 \neq t \in T$. Then, for ε small enough, the intersection*

$$\tilde{B}_\varepsilon := \tilde{U} \cap D_\varepsilon$$

(called the Milnor fibre of f) has the homology type of a bouquet of n -dimensional spheres. In particular the n -th Betti number $b_n(\tilde{B}_\varepsilon)$ (called the Milnor number m_p of p) coincides with the multiplicity of the critical point $0 \in \mathbb{C}^{n+1}$ of f as a solution to the following collection of equations

$$\frac{\partial f}{\partial x_1} = \frac{\partial f}{\partial x_2} = \cdots = \frac{\partial f}{\partial x_{n+1}} = 0 \, .$$

1.1. Milnor number from the algebraic point of view. Theorem 1.3 allows the following algebraic interpretation of the Milnor number.

Let $\mathcal{O}_0$ be the local ring of germs of holomorphic function of $\mathbb{C}^{n+1}$ at the origin. By definition of holomorphic function and the identity principle we have that $\mathcal{O}_0$ is isomorphic to the ring of convergent power series $\mathbb{C}\{x_1, \dots, x_{n+1}\}$. A germ of hypersurface singularity is defined as the Stein complex space

$$(2) \quad U_0 := \text{Spec}(\mathcal{O}_{f,0})$$

where $\mathcal{O}_{f,0} := \mathcal{O}_0/(f)$ and f is the germ represented by the series expansion of the holomorphic function (1).

Definition 1.4 (Milnor number of an i.h.s, see e.g. [16]). The *Milnor number* of the hypersurface singularity $0 \in U_0$ is defined as *the multiplicity of the critical point $0 \in \mathbb{C}^{n+1}$ of f as a solution of the system of partials of f* ([18] §7) which is

$$(3) \quad \mu_f(0) = \dim_{\mathbb{C}}(\mathcal{O}_0/J_f) = \dim_{\mathbb{C}}(\mathbb{C}\{x_1, \dots, x_{n+1}\}/J_f)$$

where $\dim_{\mathbb{C}}$ means “dimension as a $\mathbb{C}$ -vector space” and J_f is the jacobian ideal $J_f := \left(\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_{n+1}}\right)$. For shortness we will denote the Milnor number (3) by $\mu(0)$ whenever f is clear.

2. TYURINA NUMBER OF AN ISOLATED HYPERSURFACE SINGULARITY

2.1. Deformations of complex spaces. Let $\mathcal{X} \xrightarrow{x} B$ be a *flat*, surjective map of complex spaces such that B is connected and there exists a special point $0 \in B$ whose fibre $X = x^{-1}(0)$ may be singular. Then $\mathcal{X}$ is called a *deformation family* of X . If the fibre $X_b = x^{-1}(b)$ is smooth, for some $b \in B$, then X_b is called a *smoothing* of X .

Let Ω_X be the sheaf of holomorphic differential forms on X and consider the *Lichtenbaum–Schlessinger cotangent sheaves* [15] of X , $\Theta_X^i = \mathcal{E}xt^i(\Omega_X, \mathcal{O}_X)$. Then $\Theta_X^0 = \mathcal{H}om(\Omega_X, \mathcal{O}_X) =: \Theta_X$ is the “tangent” sheaf of X and Θ_X^i is supported over $\text{Sing}(X)$, for any $i > 0$. Consider the associated *local* and *global deformation objects*

$$T_X^i := H^0(X, \Theta_X^i) \quad , \quad \mathbb{T}_X^i := \text{Ext}^i(\Omega_X^1, \mathcal{O}_X) \quad , \quad i = 0, 1, 2.$$

Then by the *local to global spectral sequence* relating the global Ext and sheaf $\mathcal{E}xt$ (see [14] and [10] II, 7.3.3) we get

$$E_2^{p,q} = H^p(X, \Theta_X^q) \implies \mathbb{T}_X^{p+q}$$

giving that

$$(4) \quad \mathbb{T}_X^0 \cong T_X^0 \cong H^0(X, \Theta_X) \quad ,$$

$$(5) \quad \text{if } X \text{ is smooth then } \mathbb{T}_X^i \cong H^i(X, \Theta_X) \quad ,$$

$$(6) \quad \text{if } X \text{ is Stein then } T_X^i \cong \mathbb{T}_X^i \quad .$$

Recall that $\mathcal{X} \xrightarrow{x} B$ is called a *versal* deformation family of X if for any deformation family $(\mathcal{Y}, X) \xrightarrow{y} (C, o)$ of X there exists a map of pointed complex spaces $h : (U, o) \rightarrow (B, 0)$, defined on a neighborhood $o \in U \subset C$, such that $\mathcal{Y}|_U$ is the

pull-back of $\mathcal{X}$ by h i.e.

$$\begin{array}{ccc} \mathcal{Y}|_U = U \times_B \mathcal{X} & \longrightarrow & \mathcal{X} \\ \downarrow y & & \downarrow x \\ C \longleftarrow U & \xrightarrow{h} & B \end{array}$$

Theorem 2.1 (Douady–Grauert–Palamodov [9], [11], [21] and [22] Theorems 5.4 and 5.6). *Every compact complex space X has an effective versal deformation $\mathcal{X} \xrightarrow{x} B$ which is a proper map and a versal deformation of each of its fibers. Moreover the germ of analytic space $(B, 0)$ (the Kuranishi space of X) is isomorphic to the germ of analytic space $(q^{-1}(0), 0)$, where $q : \mathbb{T}_X^1 \rightarrow \mathbb{T}_X^2$ is a suitable holomorphic map (the obstruction map) such that $q(0) = 0$.*

In particular if $q \equiv 0$ (e.g. when $\mathbb{T}_X^2 = 0$) then $(B, 0)$ turns out to be isomorphic to the germ of a neighborhood of the origin in $\mathbb{T}_X^1$.

2.2. Deformations of an i.h.s. Let us consider the germ of i.h.s. $U_0 := \text{Spec}(\mathcal{O}_{f,0})$ as defined in (2).

Definition 2.2 (Tyurina number of an i.h.s.). The *Tyurina number* of the i.h.s. $0 \in U_0$ is

$$\tau_f(0) := \dim_{\mathbb{C}} \mathbb{T}_{U_0}^1 \stackrel{(6)}{=} \dim_{\mathbb{C}} T_{U_0}^1 = h^0(U_0, \Theta_{U_0}^1)$$

often denoted simply by $\tau(0)$ whenever f is clear. Since U_0 is Stein, the obstruction map q in Theorem 2.1 is trivial and the Tyurina number $\tau(0)$ turns out to give *the dimension of the Kuranishi space of U_0* .

Proposition 2.3 (see e.g. [25]). *If $0 \in U_0 = \text{Spec}(\mathcal{O}_{f,0})$ is the germ of an i.h.s. then $\mathbb{T}_{U_0}^1 \cong \mathcal{O}_{f,0}/J_f$ and*

$$(7) \quad \tau_f(0) = \dim_{\mathbb{C}} (\mathbb{C}\{x_1, \dots, x_{n+1}\}/I_f) \ .$$

where $I_f := (f) + J_f$. Then, recalling (3), $\tau(0) \leq \mu(0)$. In particular $\tau(0)$ gives the multiplicity of 0 as a singular point of the complex space germ U_0 .

3. MILNOR AND TYURINA NUMBERS OF A POLYNOMIAL

Let us consider the polynomial algebra $\mathbb{C}[\mathbf{x}] := \mathbb{C}[x_1, \dots, x_{n+1}]$ and let $I \subset \mathbb{C}[\mathbf{x}]$ be an ideal. If we consider the natural inclusion $\mathbb{C}[\mathbf{x}] \subset \mathbb{C}\{\mathbf{x}\} := \mathbb{C}\{x_1, \dots, x_{n+1}\}$ then we get

$$(8) \quad \dim_{\mathbb{C}} (\mathbb{C}\{\mathbf{x}\}/I \cdot \mathbb{C}\{\mathbf{x}\}) \leq \dim_{\mathbb{C}} (\mathbb{C}[\mathbf{x}]/I)$$

since the algebra $\mathbb{C}\{\mathbf{x}\}$ contains also non constant units.

Example 3.1. Let us consider the ideal $I = (x + x^2) \subset \mathbb{C}[x]$. Then $\mathbb{C}[x]/I = \langle 1, x \rangle_{\mathbb{C}}$ while $\mathbb{C}\{x\}/I \cdot \mathbb{C}\{x\} = \langle 1 \rangle_{\mathbb{C}}$, since $x \in I \cdot \mathbb{C}\{x\}$ being associated with the generator $x + x^2$ by the unit $1 + x \in \mathbb{C}\{x\}$. This means that

$$(9) \quad 1 = \dim_{\mathbb{C}} (\mathbb{C}\{x\}/I \cdot \mathbb{C}\{x\}) < \dim_{\mathbb{C}} (\mathbb{C}[x]/I) = 2 \ .$$

In particular the first equality means that, recalling Definition 1.4 and formula (3), the Milnor number of $0 \in \mathbb{C}$, as a critical point of the polynomial map $f(x) = (\frac{1}{2} + \frac{1}{3}x)x^2$, turns out to be $\mu(0) = 1$.

Let us then set the following

Definition 3.2 (Milnor and Tyurina numbers of a polynomial). Given a polynomial $f \in \mathbb{C}[\mathbf{x}]$ the following dimension

$$(10) \quad \mu(f) := \dim_{\mathbb{C}} (\mathbb{C}[\mathbf{x}]/J_f)$$

is called *the Milnor number of the polynomial f* . Analogously the dimension

$$(11) \quad \tau(f) := \dim_{\mathbb{C}} (\mathbb{C}[\mathbf{x}]/I_f)$$

where $I_f := (f) + J_f$, is called *the Tyurina number of the polynomial f* . Then clearly $\tau(f) \leq \mu(f)$.

The inequality (8) then gives that $\mu_f(p) \leq \mu(f)$ and $\tau_f(p) \leq \tau(f)$, for any point $p \in \mathbb{C}^{n+1}$ and any polynomial $f \in \mathbb{C}[\mathbf{x}]$. Moreover Milnor and Tyurina numbers of points and polynomials are related by the following

Proposition 3.3 (see e.g. [13] §A.9). *For any $f \in \mathbb{C}[x_1, \dots, x_{n+1}]$*

$$(12) \quad \begin{aligned} \mu(f) &= \sum_{p \in \mathbb{C}^{n+1}} \mu_f(p) \\ \tau(f) &= \sum_{p \in \mathbb{C}^{n+1}} \tau_f(p) \end{aligned}$$

Remark 3.4. Observe that sums on the right terms of (12) are actually finite and well-defined since $\mu(p) \neq 0$ if and only if p is a critical point of the polynomial map f and $\tau(p) \neq 0$ if and only if p is a singular point of the n -dimensional algebraic hypersurface $f^{-1}(0) \subset \mathbb{C}^{n+1}$.

Example 3.5 (Example 3.1 continued). Critical points of the polynomial map $f(x) = (\frac{1}{2} + \frac{1}{3}x)x^2$ are given by 0 and -1 . The translation $x \mapsto x - 1$ transforms f into the polynomial $g(x) = \frac{1}{3}(\frac{1}{2} + x)(x - 1)^2$ and

$$\mu_f(-1) = \mu_g(0) = \dim_{\mathbb{C}} (\mathbb{C}\{x\}/(x^2 - x)) = 1.$$

The inequality (9) then gives $2 = \mu(f) = \mu_f(0) + \mu_f(-1)$, according with the first equality in (12). Moreover 0 is the unique singular point of the 0-dimensional hypersurface $f^{-1}(0) = \{-3/2, 0\} \subset \mathbb{C}$ and

$$\begin{aligned} \tau(f) &= \dim_{\mathbb{C}} \left(\mathbb{C}[x] / \left(\left(\frac{1}{2} + \frac{1}{3}x \right) x^2, x + x^2 \right) \right) = 1 \\ \tau_f(0) &= \dim_{\mathbb{C}} \left(\mathbb{C}\{x\} / \left(\left(\frac{1}{2} + \frac{1}{3}x \right) x^2, x + x^2 \right) \right) = 1 \end{aligned}$$

according with the second equality in (12).

3.1. Milnor and Tyurina numbers of weighted homogeneous polynomials.

Let us recall that a polynomial $f \in \mathbb{C}[x_1, \dots, x_{n+1}]$ is called *weighted homogeneous (w.h.p.)* or *quasi-homogeneous* if there exist $n+1$ positive rational numbers $\mathbf{w} = (w_1, \dots, w_{n+1}) \in \mathbb{Q}^{n+1}$ such that

$$\sum_{i=1}^{n+1} w_i \alpha_i = 1$$

for any monomial $\mathbf{x}^\alpha := \prod_{i=1}^{n+1} x_i^{\alpha_i}$ appearing in f ; $\mathbf{w}$ is then called *the vector of (rational) weights of f* and the *generalized Euler formula*

$$(13) \quad f = \sum_{i=1}^{n+1} w_i x_i \frac{\partial f}{\partial x_i}$$

follows immediately for any w.h.p. $f \in \mathbb{C}[\mathbf{x}]$ admitting the same vector of weights. For any $i = 1, \dots, n+1$, let $(p_i, q_i) \in \mathbb{N}^2$ be the unique ordered couple of positive coprime integers such that p_i/q_i is the reduced fraction representing the positive rational number w_i . Calling d the least common factor of denominators $q_1, \dots, q_{n+1}$, the positive integers $d_i := dw_i$ satisfy the following *weighted homogeneity relation*

$$(14) \quad \forall \lambda \in \mathbb{C} \quad f(\lambda^{d_1} x_1, \dots, \lambda^{d_{n+1}} x_{n+1}) = \lambda^d f(x_1, \dots, x_{n+1}).$$

For this reason $\mathbf{d} = (d_1, \dots, d_{n+1})$ is called *the vector of integer weights of f* and $d = |\mathbf{d}|$ is called the *degree of f* .

Proposition 3.6. *Given a polynomial $f \in \mathbb{C}[\mathbf{x}]$ with a finite number of critical points, the following assertions are equivalent:*

- (a) f is a w.h.p.,
- (b) $\tau(f) = \mu(f)$,
- (c) $\forall p \in \mathbb{C}^{n+1} \quad \tau(p) = \mu(p)$.

In particular (c) implies that the set of critical points of f coincides with the set of singular points of $f^{-1}(0)$. Moreover (a) means actually that the origin $0 \in \mathbb{C}^{n+1}$ is the unique possible critical point of f and then the unique possible singular point of $f^{-1}(0)$. Therefore (b) and (c) give that

$$(15) \quad \mu(0) = \mu(f) = \tau(f) = \tau(0).$$

Proof. (a) $\Rightarrow$ (c). The generalized Euler formula (13) implies that $I_f = J_f$ and (c) follows immediately by (3) and (7). In particular if $\mathbf{x} = (x_1, \dots, x_{n+1})$ is a critical point of f , then (13) and (14) give that $(\lambda^{d_1} x_1, \dots, \lambda^{d_{n+1}} x_{n+1})$ is a critical point of f for any complex number λ . Then $\mathbf{x} = 0$ since f admits at most a finite number of critical points, meaning that f admits at most the origin as a critical point.

(c) $\Rightarrow$ (b). This follows immediately by Proposition 3.3.

(b) $\Rightarrow$ (a). Since $J_f \subseteq I_f$ then we get the natural surjective map of $\mathbb{C}$ -algebras

$$\mathbb{C}[\mathbf{x}]/J_f \longrightarrow \mathbb{C}[\mathbf{x}]/I_f$$

The hypothesis $\tau(f) = \mu(f)$ implies that it is also injective which suffices to show that $J_f = I_f$. Hence $f \in J_f$ and a famous result by K. Saito [24] allows to conclude that f is a w.h.p. $\square$

Definition 3.7 (Weighted homogeneous singularity - w.h.s.). A n -dimensional i.h.s. $0 \in U_0 = \text{Spec } \mathcal{O}_{f,0}$ is called *weighted homogeneous* (or *quasi-homogeneous*) if there exists a w.h.p. $F \in \mathbb{C}[x_1, \dots, x_{n+1}]$ such that $U_0 \cong \text{Spec } \mathcal{O}_{F,0}$, as germs of complex spaces.

Remark 3.8. Definition 3.7 is equivalent to require that there exists an automorphism ϕ^* of $\mathcal{O}_0 = \mathbb{C}\{\mathbf{x}\}$, induced by a biholomorphic local coordinates change $(\mathbb{C}^{n+1}, 0) \xrightarrow{\phi} (\mathbb{C}^{n+1}, 0)$, such that $\phi^*(f) := f \circ \phi = F$ (see [12] Lemma 2.13).

Proposition 3.9 (Characterization of a w.h.s.). $0 \in U_0 = \text{Spec } \mathcal{O}_{f,0}$ is a w.h.s. if and only if $\tau_f(0) = \mu_f(0)$.

Proof. The statement follows immediately by Proposition 3.6, keeping in mind Remark 3.8 and observing that the jacobian ideals J_f and J_F can be obtained each other by multiplying the jacobian matrix of the coordinate change ϕ , which is clearly invertible in a neighborhood of 0. $\square$

3.2. An example: weighted homogeneous cDV singularities. An example of an isolated hypersurface singularity is given by a *compound Du Val* (cDV) singularity which is a 3-fold point p such that, for a hyperplane section H through p , $p \in H$ is a Du Val surface singularity i.e. an A–D–E singular point (see [23], §0 and §2, and [6], chapter III). Then a cDV point p is a germ of hypersurface singularity $0 \in U_0 := \text{Spec}(\mathcal{O}_{f,0})$, where f is the polynomial

$$(16) \quad f(x, y, z, t) := x^2 + q(y, z) + t g(x, y, z, t) ,$$

such that $g(x, y, z, t)$ is a generic element of the maximal ideal $\mathfrak{m}_0 := (x, y, z, t) \subset \mathbb{C}[x, y, z, t]$ and

$$(17) \quad \begin{aligned} A_n &: q(y, z) := y^2 + z^{n+1} \quad \text{for } n \geq 1 \\ D_n &: q(y, z) := y^2 z + z^{n-1} \quad \text{for } n \geq 4 \\ E_6 &: q(y, z) := y^3 + z^4 \\ E_7 &: q(y, z) := y^3 + yz^3 \\ E_8 &: q(y, z) := y^3 + z^5 \end{aligned}$$

In particular if

$$(18) \quad g(x, y, z, t) = t$$

then $f = 0$ in (16) is said to define an *Arnol'd simple (threefold) singularity* ([3], [5] §15 and in particular [4] §I.2.3) denoted by $A_n, D_n, E_{6,7,8}$, respectively.

The index $(n, 6, 7, 8)$ turns out to be the Milnor number of the surface Du Val singularity $0 \in U_0 \cap \{t = 0\}$ or equivalently its Tyurina number, since a Du Val singular point always admits a weighted homogeneous local equation. When a cDV point is defined by a weighted homogeneous polynomial f , a classical result of J. Milnor and P. Orlik allows to compare this index with its Milnor (and then Tyurina) number. In particular we get

$$(19) \quad \begin{aligned} w(x) &= 1/2 \\ w(y) &= \begin{cases} 1/2 & \text{if } p \text{ is } cA_n, \\ (n-2)/(2n-2) & \text{if } p \text{ is } cD_n, \\ 1/3 & \text{if } p \text{ is } cE_{6,7,8}. \end{cases} \\ w(z) &= \begin{cases} 1/(n+1) & \text{if } p \text{ is } cA_n, \\ 1/(n-1) & \text{if } p \text{ is } cD_n, \\ 1/4 & \text{if } p \text{ is } cE_6, \\ 2/9 & \text{if } p \text{ is } cE_7, \\ 1/5 & \text{if } p \text{ is } cE_8. \end{cases} \end{aligned}$$

Theorem 3.10 (Milnor–Orlik [19], Thm. 1). *Let $f(x_1, \dots, x_{n+1})$ be a w.h.p., with rational weights $w_1, \dots, w_{n+1}$, admitting an isolated critical point at the origin. Then the Milnor number of the origin is given by*

$$\mu(0) = (w_1^{-1} - 1) (w_2^{-1} - 1) \cdots (w_{n+1}^{-1} - 1) .$$

By putting weights (19) in the previous Milnor–Orlik formula we get the following

Corollary 3.11. *Let $0 \in U_0$ be a w.h. cDV point of index n . Then*

$$\tau(0) = \mu(0) = n(w(t)^{-1} - 1) .$$

In particular for Arnol'd simple singularities we get $n = \tau(0) = \mu(0)$, as can also be directly checked by the definition.

3.3. The algebraic computation via Gröbner basis. Let us consider an ideal $I \subset \mathbb{C}[\mathbf{x}]$ and let $L(I)$ denote the ideal generated by *leading monomials*, with respect to a fixed *monomial order*, of elements in I . It is a well known fact (see e.g. [7] §5.3) that the following are isomorphisms of $\mathbb{C}$ -vector spaces

$$(20) \quad \mathbb{C}[x_1, \dots, x_{n+1}]/I \xrightarrow{\cong} \mathbb{C}[x_1, \dots, x_{n+1}]/L(I) \xrightarrow{\cong} \langle M \setminus L(I) \rangle_{\mathbb{C}}$$

where M is the set (actually a multiplicative monoid) of all monomials $\mathbf{x}^\alpha$. Consider a polynomial $f \in \mathbb{C}[\mathbf{x}]$. By Definition 3.2 and isomorphisms (20), the computation of $\mu(f)$ and $\tau(f)$ reduces to calculate

$$\dim_{\mathbb{C}} (\mathbb{C}[x_1, \dots, x_{n+1}]/L(I)) = |M \setminus L(I)|$$

where $L(I)$ is the ideal generated by *leading monomials*, with respect to a fixed monomial order, of elements in the ideal $I \subset \mathbb{C}[x_1, \dots, x_{n+1}]$ which is either the jacobian ideal J_f or the ideal $I_f = (f) + J_f$, respectively. The point is then *determining a Gröbner basis of I w.r.t. the fixed monomial order*, which can be realized e.g. by the **Groebner Package** of MAPLE.

Remark 3.12. The MAPLE computation of Milnor and Tyurina numbers of polynomials is realized by procedures **PolyMilnor** and **PolyTyurina**, whose concrete description is postponed to section 5. Their usefulness is clarified by Proposition 3.6 and in particular by equations (15). In fact in the case of a w.h.p. f admitting an isolated critical point in $0 \in \mathbb{C}^{n+1}$ there is no need of working with power series (and then with *local monomial orders*) to determine $\mu(0)$ and $\tau(0)$. Since the Buchberger algorithm implemented with MAPLE turns out to be more efficient when running by usual term orders, the use of *global* procedures **PolyMilnor** and **PolyTyurina** has to be preferred to the use of their *local* counterparts **Milnor** and **Tyurina**, when possible.

4. MONOMIAL ORDERING

First of all let us recall what is usually meant by a monomial order. For more details the interested reader is remanded e.g. to [7] §2.2, [13] §9 and [8] §1.

Let M be the multiplicative monoid of monomials $\mathbf{x}^\alpha = \prod_{i=1}^{n+1} x_i^{\alpha_i}$: clearly $\log : M \xrightarrow{\cong} \mathbb{N}^{n+1}$. A (*global*) *monomial order* on $\mathbb{C}[\mathbf{x}]$ is a *total order relation* $\leq$ on M which is

- (i) *multiplicative* i.e. $\forall \alpha, \beta, \gamma \in \mathbb{N}^{n+1} \quad \mathbf{x}^\alpha \leq \mathbf{x}^\beta \Rightarrow \mathbf{x}^\alpha \cdot \mathbf{x}^\gamma \leq \mathbf{x}^\beta \cdot \mathbf{x}^\gamma$,
- (ii) *a well-ordering* i.e. every nonempty subset of M has a *smallest element*.

Since $\mathbb{C}[\mathbf{x}]$ is a noetherian ring a multiplicative total order on M is a well-ordering if and only if

- (ii') $\forall i = 1, \dots, n+1 \quad 1 < x_i$.

Definition 4.1 (Local and global monomial orders, [13] Definition 1.2.4). In the following a m.o. on $\mathbb{C}[\mathbf{x}]$ will denote simply a total order relation $\leq$ on M which is multiplicative i.e. satisfying (i). A m.o. will be called *global (g.m.o.)* if also (ii), or equivalently (ii'), is satisfied. Moreover a m.o. will be called *local (l.m.o.)* if

(ii'') $\forall i = 1, \dots, n+1 \quad x_i < 1$.

4.1. Localizations in $\mathbb{C}[\mathbf{x}]$ and rings implemented by monomial orders.

Given a m.o. $\leq$ on M consider the following subset of $\mathbb{C}[\mathbf{x}]$

$$S := \{f \in \mathbb{C}[\mathbf{x}] \mid L(f) \in \mathbb{C} \setminus \{0\}\}$$

where $L(f)$ is the *leading monomial* of f w.r.t. $\leq$. Since S is a multiplicative subset of $\mathbb{C}[\mathbf{x}]$ we can consider the localization $S^{-1}\mathbb{C}[\mathbf{x}]$. Then (ii') and (ii'') give immediately the following

Proposition 4.2.

$$S^{-1}\mathbb{C}[\mathbf{x}] = \mathbb{C}[\mathbf{x}] \Leftrightarrow \leq \text{ is a g.m.o.}$$

$$S^{-1}\mathbb{C}[\mathbf{x}] = \mathbb{C}[\mathbf{x}]_{(\mathbf{x})} \Leftrightarrow \leq \text{ is a l.m.o.}$$

where $\mathbb{C}[\mathbf{x}]_{(\mathbf{x})}$ is the localization of $\mathbb{C}[\mathbf{x}]$ at the maximal ideal $(\mathbf{x}) \subset \mathbb{C}[\mathbf{x}]$.

By Taylor power series expansion of locally holomorphic functions, there is a natural inclusion $\mathbb{C}[\mathbf{x}]_{(\mathbf{x})} \subset \mathbb{C}\{\mathbf{x}\}$ giving the following commutative diagram, for every ideal $I \subset \mathbb{C}[\mathbf{x}]_{(\mathbf{x})}$:

$$\begin{array}{ccc} \mathbb{C}[\mathbf{x}]_{(\mathbf{x})} & \hookrightarrow & \mathbb{C}\{\mathbf{x}\} \\ \downarrow & & \downarrow \\ \mathbb{C}[\mathbf{x}]_{(\mathbf{x})}/I & \hookrightarrow & \mathbb{C}\{\mathbf{x}\}/I \cdot \mathbb{C}\{\mathbf{x}\} \end{array}$$

Proposition 4.3 (for a proof see e.g. [8] Proposition 9.4). *If $\dim_{\mathbb{C}} (\mathbb{C}[\mathbf{x}]_{(\mathbf{x})}/I)$ is finite then the inclusion $\mathbb{C}[\mathbf{x}]_{(\mathbf{x})}/I \subset \mathbb{C}\{\mathbf{x}\}/I \cdot \mathbb{C}\{\mathbf{x}\}$ is an isomorphism of $\mathbb{C}$ -algebras. In particular both the underlying vector spaces have the same dimension.*

Theorem 4.4 ([8] Theorem 9.29). *Let $\leq$ be a m.o. on $\mathbb{C}[\mathbf{x}]$ and $I \subset S^{-1}\mathbb{C}[\mathbf{x}]$ be an ideal. Then*

$$\dim_{\mathbb{C}} (S^{-1}\mathbb{C}[\mathbf{x}]/I) = \dim_{\mathbb{C}} (\mathbb{C}[\mathbf{x}]/L(I))$$

where $L(I)$ is the ideal generated by leading monomials, with respect to $\leq$, of polynomials in I . In particular if it is finite then $M \setminus L(I)$ represents a basis of the vector space $S^{-1}\mathbb{C}[\mathbf{x}]$.

Corollary 4.5. *Let $f \in \mathbb{C}[\mathbf{x}]$ admit an isolated critical point at $0 \in \mathbb{C}^{n+1}$, $\leq$ be a m.o. on $\mathbb{C}[\mathbf{x}]$ and $I = J_f$ (resp. $I = I_f$). Then*

$$\dim_{\mathbb{C}} (\mathbb{C}[\mathbf{x}]/L(I)) = \begin{cases} \mu(f) \text{ (resp. } \tau(f)) & \text{if } \leq \text{ is global} \\ \mu_f(0) \text{ (resp. } \tau_f(0)) & \text{if } \leq \text{ is local} \end{cases}$$

Remark 4.6. The point is then determining a *standard basis* of I w.r.t. the fixed m.o. $\leq$. If the latter is a *global* one, a standard basis is a usual Gröbner basis which is obtained by applying the Buchberger algorithm. In MAPLE this is implemented by the **Groebner Package**.

On the other hand, if $\leq$ is a *local* m.o. the Buchberger algorithm does no more work. In fact $\leq$ is no more a well-ordering and the division algorithm employed by the Buchberger algorithm for determining *normal forms* of *S-polynomials* may do not terminate. This problem can be dodged by means of a *weak normal form algorithm* (*weakNF*), firstly due to F. Mora [20], and of a *standard basis algorithm* (*SB*) which replaces the Buchberger algorithm. This is precisely what has been

implemented in SINGULAR since 1990 (see [13] §1.7 and references thereof). The aim of the following section 5 is to present a MAPLE implementation of *weakNF* and *SB* algorithms to yield a procedure computing Milnor and Tyurina numbers of points.

5. MAPLE SUBROUTINES IN DETAIL

The present section is devoted to present and describe in detail MAPLE subroutines computing Milnor and Tyurina numbers of critical points of a polynomial f . They are available as MAPLE 12 files `.mw` at [1]. They are composed by several procedures, the most important of which are the following:

- **weakNF** which is the MAPLE implementation of the weak normal form Algorithm 1.7.6 in [13]:

Input : $f \in \mathbb{C}[\mathbf{x}]$, G a finite subset in $\mathbb{C}[\mathbf{x}]$,
 $\text{variables} \subseteq \mathbf{x} = \{x_1, \dots, x_{n+1}\}$,
 STO = a MAPLE type `ShortMonomialOrder` expression
Output : $h \in (\mathbb{C}[\mathbf{x} \setminus \text{variables}])[\text{variables}]$
which is a polynomial weak normal form of f w.r.t. G ;

- **SB** which is the MAPLE implementation of the standard basis Algorithm 1.7.1 in [13]:

Input : G a finite subset in $\mathbb{C}[\mathbf{x}]$, **weakNF**,
Output : a standard basis S of $I := (G) \subset \mathbb{C}[\mathbf{x}]$;

- **Milnor** which is the procedure computing the Milnor number of an isolated critical point:

Input : $F \in \mathbb{C}[\mathbf{x}]$, **SB**,
Optional: **variables**, by default assigned by `indets(F)`
Optional: a l.m.o., by default assigned by `tdeg_min(variables)`
Output : A standard basis G of $J_F \subset (\mathbb{C}[\mathbf{x} \setminus \text{variables}])[\text{variables}]$, $L(G)$,
 $M \setminus L(J_F)$, $\mu_F(0)$;

- **PolyMilnor** which is the procedure computing the Milnor number of a polynomial:

Input : $F \in \mathbb{C}[\mathbf{x}]$, **Basis** in the **Groebner** package,
Optional: **variables**, by default assigned by `indets(F)`
Optional: a g.m.o., by default assigned by `tdeg(variables)`
Output : A Gröbner basis G of $J_F \subset (\mathbb{C}[\mathbf{x} \setminus \text{variables}])[\text{variables}]$, $L(G)$,
 $M \setminus L(J_F)$, $\mu(F)$;

- **Tyurina** which is the procedure computing the Tyurina number of an isolated singular point:

Input : $F \in \mathbb{C}[\mathbf{x}]$, **SB**,

Optional: **variables**, by default assigned by **indets(F)**

Optional: a l.m.o., by default assigned by **tdeg_min(variables)**

Output : A standard basis G of $I_F \subset (\mathbb{C}[\mathbf{x} \setminus \text{variables}])[\text{variables}]$, $L(G)$,
 $M \setminus L(I_F)$, $\tau_F(0)$;

- **PolyTyurina** which is the procedure computing the Tyurina number of a polynomial:

Input : $F \in \mathbb{C}[\mathbf{x}]$, **Basis** in the **Groebner** package,

Optional: **variables**, by default assigned by **indets(F)**

Optional: a g.m.o., by default assigned by **tdeg(variables)**

Output : A Gröbner basis G of $I_F \subset (\mathbb{C}[\mathbf{x} \setminus \text{variables}])[\text{variables}]$, $L(G)$,
 $M \setminus L(I_F)$, $\tau(F)$.

5.1. **The subroutines.** Preambles to introduce the useful MAPLE packages:

```
> with(Groebner):
> with(PolynomialIdeals):
> with(Ore_algebra):
```

A first control procedure :

```
> localorglobal := proc (ST0, variables)
> A:= poly_algebra(op(variables)); TP := MonomialOrder(A, ST0);
> nuu:= 1; muu := 1;
> for i to nops(variables) do if TestOrder(1,variables[i], TP)
> then nuu := 0 else muu := 0 end if end do;
> if nuu = 1 then Lo else if muu = 1 then G1 else Mi
> end if end if end proc:
```

Actually **localorglobal** is not an essential procedure in the present routine: its meaning is simply that of giving a feedback about what kind of m.o. the user is employing and stopping the procedure running with a wrong term order: in fact **Milnor** and **Tyurina** may give wrong output when running with a g.m.o.; on the other hand **PolyMilnor** and **PolyTyurina** may not terminate when running with a l.m.o..

5.1.1. *Implementing local monomial orders.* The following three procedures give the core of the *MAPLE implementation of l.m.o.'s* for determining standard basis of ideals in $\mathbb{C}[\mathbf{x}]_{(\mathbf{x})}$. The first procedure introduces the *ecart* concept which is the main ingredient in the Mora algorithm **weakNF**. It is defined following [13] Definition 1.7.5:

```
> ecart := proc (f, variables, ST0)
> degree(f, variables)-degree(LeadingMonomial(f, ST0), variables)
> end proc:
```

Then we give the Mora algorithm for determining a weak normal form of a polynomial $f \in \mathbb{C}[\mathbf{x}]$ w.r.t. a finite subset of polynomials $G \subset \mathbb{C}[\mathbf{x}]$ (see [13] Algorithm 1.7.6 and [8] Algorithm 9.22):

```

> weakNF := proc (f, G, variables, ST0)
>   h := f;
>   TT := G;
>   TTh := {}; for i to nops(TT) do
>     if divide(LeadingMonomial(h, ST0), LeadingMonomial(TT[i], ST0))
>     then TTh := {TT[i], op(TTh)}
>     end if end do;
>   while (h <> 0 and TTh <> {}) do
>     L := [op(TTh)];
>     L1 := sort(L, proc (t1, t2) options operator, arrow;
>       ecart(t1, variables, ST0) <= ecart(t2, variables, ST0) end proc);
>     g := L1[1];
>     if ecart(h, variables, ST0) < ecart(g, variables, ST0)
>     then TT := {h, op(TT)} end if;
>     h := SPolynomial(h, g, ST0);
>     TTh := {};
>     for i to nops(TT) do
>       if divide(LeadingMonomial(h, ST0), LeadingMonomial(TT[i], ST0))
>       then TTh := {TT[i], op(TTh)} end if end do end do;
>   h end proc

```

At last the standard basis procedure giving the analogue of Buchberger algorithm with l.m.o.'s:

```

> SB := proc (G, variables, ST0)
>   S := G;
>   P := {seq(seq({G[i], G[j]}, j=i+1..nops(G)), i=1..nops(G))};
>   while P <> {} do
>     P1 := P[1];
>     P := 'minus'(P, {P1});
>     h := weakNF(SPolynomial(P1[1], P1[2], ST0), S, variables, ST0);
>     if h <> 0 then P := {seq({h, S[i]}, i = 1 .. nops(S)), op(P)};
>     S := {h, op(S)} end if end do;
>   S end proc:

```

5.1.2. *Computing Milnor and Tyurina numbers.* We are now in a position to introduce the procedures computing Milnor and Tyurina numbers of critical points of a polynomial $F \in \mathbb{C}[\mathbf{x}]$. They actually give some more output. Precisely, saying $I = J_F$ (resp. $I = I_F$) Milnor (resp. Tyurina) returns:

- a standard basis G of the ideal $I \cdot \mathbb{C}[\mathbf{x}]_{(\mathbf{x})} \subset \mathbb{C}[\mathbf{x}]_{(\mathbf{x})}$,
- the leading monomial basis $L(G)$ w.r.t. a fixed m.o.,
- $M \setminus L(I)$ representing $(\text{ mod } I \cdot \mathbb{C}[\mathbf{x}])$ a basis of the quotient vector space $\mathbb{C}[\mathbf{x}]_{(\mathbf{x})} / I \cdot \mathbb{C}[\mathbf{x}]_{(\mathbf{x})}$
- and its dimension over $\mathbb{C}$, which is $\mu_F(0)$ (resp. $\tau_F(0)$).

Each of the previous output may be listed separately by means of the sub-procedures `MilnorGroebnerBasis`, `MilnorLT`, `MilnorBasis` and `MilnorNumber` (respectively, `TyurinaGroebnerBasis`, `TyurinaLT`, `TyurinaBasis` and `TyurinaNumber`).

The Milnor procedure:

```

> Milnor := proc (F, variables::set:=indets(F),
> U::anything:=tdeg_min(op(variables)))
> if type(U,ShortMonomialOrder) = false then error
> "invalid input: your ShortMonomialOrder is not well defined"
> else if localorglobal(U, variables) <> Lo then error
> "invalid input: your Short Monomial Order should be LOCAL"
> else r := nops(variables);
> J := [seq(diff(F, variables[i]), i = 1..r)];
> G := SB(J,variables, U);
> Ini := [seq(LeadingMonomial(G[i], U), i = 1..nops(G))];
> massimo:=r*max(seq(degree(Ini[i],variables),i =1..nops(Ini)));
> if IsZeroDimensional(<(op(Ini))>,{op(variables)}) = false then
> error "the given critical point is not isolated"
> else
> L := sort([op(subs(uq = 1, convert(map(expand,
> series(1/(product(1-variables[k]*uq, k = 1 .. r)),
> uq, massimo+1)), polynom))],
> proc (t1, t2) options operator, arrow;
> TestOrder(t1, t2, U) end proc);
> M := [];
> for j to nops(L) do up := 1;
> for k to nops(Ini) do
> if divide(L[j], Ini[k]) then up := 0
> end if end do;
> if up = 1 then M := [op(M), L[j]]
> end if end do;
> [G,Ini, M, nops(M)]
> end if end if end if end if end proc:

```

The associated sub-procedures:

```

> MilnorGroebnerBasis:=proc(F,variables::set:=indets(F),
> T::anything:=tdeg_min(op(variables)))
> Milnor(F,variables,T)[1]
> end proc:

```

```

> MilnorLT:=proc(F,variables::set:=indets(F),
> T::anything:=tdeg_min(op(variables)))
> Milnor(F, variables, T)[2]
> end proc:

```

```

> MilnorBasis:=proc(F,variables::set:=indets(F),
> T::anything:=tdeg_min(op(variables)))
> Milnor(F, variables,T)[3]
> end proc:

```

```

> MilnorNumber:=proc(F,variables::set:=indets(F),
> T::anything:=tdeg_min(op(variables)))
> Milnor(F,variables,T)[4] end proc:

```

The Tyurina procedure:

```

> Tyurina := proc(F,variables::set:=indets(F),
> U::anything := tdeg_min(op(variables)))
> if type(U,ShortMonomialOrder)=false then error
> "invalid input: your ShortMonomialOrder is not well defined":
> else if localorglobal(U, variables)<>Lo then error
> "invalid input: your Short Monomial Order should be LOCAL":
> else r := nops(variables):
> K := [F, seq(diff(F, variables[i]), i =1 .. r)]:
> H := SB(K, variables,U);
> Ini := [seq(LeadingMonomial(H[i], U), i = 1..nops(H))];
> massimo:=r*max(seq(degree(Ini[i]),i = 1..nops(Ini)));
> if IsZeroDimensional(<(op(Ini))>,{op(variables)}) = false then
> error "the given singular point is not isolated"
> else
> L := sort([op(subs(uq = 1, convert(map(expand,
> series(1/(product(1-variables[k]*uq, k = 1 .. r)),
> uq, massimo+1)), polynom))],
> proc (t1, t2) options operator, arrow;
> TestOrder(t1, t2, U) end proc);
> M := [];
> for j to nops(L) do up :=1;
> for k to nops(Ini) do
> if divide(L[j], Ini[k]) then up := 0
> end if end do;
> if up = 1 then M := [op(M), L[j]]
> end if end do;
> [H,Ini,M,nops(M)] end if end if end if end if end proc:

```

The associated sub-procedures:

```

> TyurinaGroebnerBasis := proc(F,variables::set := indets(F),
> T::anything :=tdeg_min(op(variables)))
> Tyurina(F,variables,T)[1]
> end proc:

> TyurinaLT := proc (F,variables::set :=indets(F),
> T::anything := tdeg_min(op(variables)))
> Tyurina(F,variables,T)[2]
> end proc:

> TyurinaBasis := proc (F, variables::set:= indets(F),
> T::anything := tdeg_min(op(variables)))
> Tyurina(F,variables,T)[3]
> end proc:

> TyurinaNumber := proc (F, variables::set:= indets(F),
> T::anything := tdeg_min(op(variables)))
> Tyurina(F,variables, T)[4]
> end proc:

```

Ultimately the following procedures allows to compute Milnor and Tyurina numbers $\mu(F)$ and $\tau(F)$ of a polynomial $F \in \mathbb{C}[\mathbf{x}]$. They give the same output of `Milnor` and `Tyurina` but for an ideal $I \subset \mathbb{C}[\mathbf{x}]$, since they works with a g.m.o.. As before each output may be listed separately by means of analogous sub-procedures.

The procedure computing $\mu(F)$:

```

> PolyMilnor := proc (F,variables::set:=indets(F),
> U::anything := tdeg(op(variables)))
> if type(U, ShortMonomialOrder) = false then error
> "invalid input: your ShortMonomialOrder is not well defined":
> else if localorglobal(U, variables) <> G1 then error
> "invalid input: your Short Monomial Order should be GLOBAL":
> else r := nops(variables);
> J := [seq(diff(F, variables[i]), i = 1 .. r)];
> G := Basis(J, U);
> Ini := [seq(LeadingMonomial(G[i], U), i = 1 .. nops(G))];
> massimo:=r*max(seq(degree(Ini[i]),i=1..nops(Ini)));
> if IsZeroDimensional(<(op(Ini))>, {op(variables)}) = false then
> error "there are non isolated critical points"
> else
> L := sort([op(subs(uq = 1, convert(map(expand,
> series(1/(product(1-variables[k]*uq, k = 1 .. r)),
> uq, massimo+1)), polynom))],
> proc (t1, t2) options operator, arrow;
> TestOrder(t1, t2, U) end proc);
> M := [];
> for j to nops(L) do up := 1;
> for k to nops(Ini) do
> if divide(L[j], Ini[k]) then up := 0
> end if end do;
> if up = 1 then M := [op(M), L[j]]
> end if end do;
> [G, Ini, M, nops(M)]
> end if end if end if end if end proc:

```

The associated sub-procedures:

```

> PolyMilnorGroebnerBasis := proc (F,variables::set := indets(F),
> T::anything :=tdeg(op(variables)))
> PolyMilnor(F,variables,T)[1]
> end proc:

> PolyMilnorLT := proc (F, variables::set:= indets(F),
> T::anything := tdeg(op(variables)))
> PolyMilnor(F, variables, T)[2]
> end proc:

> PolyMilnorBasis := proc (F,variables::set:= indets(F),
> T::anything :=tdeg(op(variables)))
> PolyMilnor(F,variables,T)[3]
> end proc:

> PolyMilnorNumber:=proc(F,variables::set:= indets(F),
> T::anything :=tdeg(op(variables)))
> PolyMilnor(F,variables,T)[4]
> end proc:

```

The procedure computing $\tau(F)$:

```

> PolyTyurina := proc (F, variables::set := indets(F),
> U::anything := tdeg(op(variables)))
> if type(U, ShortMonomialOrder) = false then error
> "invalid input: your ShortMonomialOrder is not well defined":
> else if localorglobal(U, variables) <> G1 then error
> "invalid input: your Short Monomial Order should be GLOBAL"
> else r := nops(variables);
> K := [F, seq(diff(F, variables[i]), i = 1 .. r)];
> H := Basis(K, U);
> Ini := [seq(LeadingMonomial(H[i], U), i = 1 .. nops(H))];
> massimo:=r*max(seq(degree(Ini[i]), i=1..nops(Ini)));
> if IsZeroDimensional(<(op(Ini))>, {op(variables)}) = false then
> error "there are non isolated singular points"
> else
> L := sort([op(subs(uq = 1, convert(map(expand,
> series(1/(product(1-variables[k]*uq, k = 1 .. r)),
> uq, massimo+1)), polynom))],
> proc (t1, t2) options operator, arrow;
> TestOrder(t1, t2, U) end proc);
> M := [];
> for j to nops(L) do up := 1;
> for k to nops(Ini) do
> if divide(L[j], Ini[k]) then up := 0
> end if end do;
> if up = 1 then M := [op(M), L[j]]
> end if end do;
> [H, Ini, M, nops(M)]
> end if end if end if end if end proc:

```

The associated sub-procedures:

```

> PolyTyurinaGroebnerBasis := proc(F, variables::set := indets(F),
> T::anything := tdeg(op(variables)))
> PolyTyurina(F, variables, T)[1]
> end proc:

> PolyTyurinaLT := proc (F, variables::set := indets(F),
> T::anything := tdeg(op(variables)))
> PolyTyurina(F, variables, T)[2]
> end proc:

> PolyTyurinaBasis := proc (F, variables,
> T::anything := tdeg(op(variables)))
> PolyTyurina(F, variables, T)[3]
> end proc:

> PolyTyurinaNumber := proc (F, variables::set := indets(F),
> T::anything := tdeg(op(variables)))
> PolyTyurina(F, variables, T)[4]
> end proc:

```

5.2. Some user friendly examples. Once implemented the subroutine 5.1 needs quite simple and minimal commands to work. As a first example let us start, for comparison, by a problem already studied by using SINGULAR in [12] Example 2.7.2(2).

Example 5.1. Let us study critical points of $F(x, y) := x^5 + y^5 + x^2y^2$ and singularities of $F^{-1}(0)$. Hence we have to type:

```

> F:=x^5+y^5+x^2*y^2:

```

Let us find, at first, the critical points of F , by solving the algebraic system of partial derivatives:

```
> solve({diff(F, x), diff(F, y)}, [x,y]);
[[x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0],
[x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0], [x = -2/5, y = -2/5],
[x = 2/5 - 2/5 (RootOf (-Z^4 - Z^3 + Z^2 - Z + 1, label = L2))^3
+ 2/5 (RootOf (-Z^4 - Z^3 + Z^2 - Z + 1, label = L2))^2
- 2/5 RootOf (-Z^4 - Z^3 + Z^2 - Z + 1, label = L2),
y = 2/5 RootOf (-Z^4 - Z^3 + Z^2 - Z + 1, label = L2)]]
```

Then F admits 6 critical points: the repetition of the solution in the origin means that this point is a multiple solution. Then we have to expect $\mu(0) > 0$.

Singular points of $F^{-1}(0)$ are given by:

```
> solve({F, diff(F, x), diff(F, y)}, [x,y])
[[x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0],
[x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0],
[x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0],
[x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0],
[x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0],
[x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0], [x = 0, y = 0],
[x = 0, y = 0]]
```

Then $F^{-1}(0)$ has a unique singular point in the origin. Therefore, by the second formula in (12), $\tau(F) = \tau(0)$ meaning that **TyurinaNumber** and **PolyTyurinaNumber** give the same number. To get this number one simply have to type either

```
> TyurinaNumber(F)
10
```

or

```
> PolyTyurinaNumber(F)
10
```

- *In general, if $F^{-1}(0)$ admits a unique singular point the procedure **PolyTyurina** has to be preferred, since it turns out to be more efficient.*

To compute Milnor numbers let us start by $\mu(F)$, by typing

```
> PolyMilnorNumber(F)
16
```

Since $\tau(0) \leq \mu(0)$ and F admits 6 critical points, by (12) we have to expect $10 \leq \mu(0) \leq 11$. Furthermore the 5 critical points different from the origin can be exchanged each other under the action of the order 5 cyclic group

$$\left\langle \left(\begin{pmatrix} -\epsilon^3 & 0 \\ 0 & \epsilon^2 \end{pmatrix} \right)^i \mid \epsilon^5 + 1 = 0, 1 \leq i \leq 5 \right\rangle \subset \text{GL}(2, \mathbb{C})$$

which is also a subgroup of $\text{Aut}(F)$. Then they cannot assume Milnor number greater than 1, giving $\mu(0) = 11$. In fact:

```
> MilnorNumber(F)
```

11

Then in this case, the use of `solve`, `PolyTyurina` and `PolyMilnor`, may avoid to employ `Tyurina` and `Milnor` which turn out to be in general less efficient procedures.

Example 5.2 (w.h. polynomials). What observed at the end of the previous Example 5.1 is obviously true for a w.h.p., after Proposition 3.6. Let us in fact consider the E_6 3-dimensional singularity $0 \in F^{-1}(0)$ where

$$F(x, y, z, t) = x^2 + y^3 + z^4 + t^2.$$

By Corollary 3.11, since $w(t) = 1/2$, one has to expect $\tau(0) = \mu(0) = 6$. This fact can be checked by the quicker procedure `PolyMilnor`:

```
> F:=x^2+y^3+z^4+t^2:
> PolyMilnorNumber(F);
```

6

Example 5.3 (Non-isolated singularities). All the procedures presented in 5.1.2 stop, giving an error message, if the considered polynomial admits *non-isolated singularities*. Consider, in fact, $F = x^2z^2 + y^2z^2 + x^2y^2$ admitting the union of the three coordinate axes as the locus of critical (and singular, since F is homogeneous) points as can easily checked by typing:

```
> F:=x^2*z^2+y^2*z^2+x^2*y^2:
> solve({diff(F, x), diff(F, y), diff(F,z)}, [x, y, z]);
```

```
[[x = 0, y = y, z = 0], [x = 0, y = y, z = 0], [x = x, y = 0, z = 0], [x = 0, y = 0, z = z]]
```

Then:

```
> PolyMilnorNumber(F);
```

Error, (in PolyMilnor) there are non isolated critical points

The user will obtain the similar error messages by running any of the other procedures.

Remark 5.4 (Be careful with variables!). The second input of any procedure in 5.1.2 is the set `variables` of variables one wants to work with. It is an *optional input* meaning that by default `variables` is assumed to be the set `indets(F)` of variables appearing in the polynomial F . This means that if the user is interesting in consider the cylinder $F^{-1}(0)$ where $F : \mathbb{C}^3 \rightarrow \mathbb{C}$ is the polynomial map $F(x, y, z) = y^2 - x(x-1)(x-2)$ then he have to type:

```
> F := y^2-x*(x-1)*(x-2):
> PolyMilnor(F, {x, y, z});
```

Error, (in PolyMilnor) there are non isolated critical points

which is right since F do not admit isolated critical points as can be checked by:

```
> solve({diff(F, x), diff(F, y), diff(F,z)}, [x,y,z]);
```

```
[[x = RootOf (3_Z^2 - 6_Z + 2), y = 0, z = z]]
```

By the way, if the set of variables is not specified then by default it is assumed to be $\{x, y\}$, meaning that F is considered as a polynomial map from $\mathbb{C}^2$ to $\mathbb{C}$. In this case F admits only the two isolated critical points

```
> solve(diff(F, x), diff(F, y), [x,y])
[[x = RootOf (3 _Z^2 - 6 _Z + 2, label = _L1) , y = 0]]
```

In fact

```
> PolyMilnor(F);
[[y, 3 x^2 - 6 x + 2], [y, x^2], [1, x], 2]
```

Read the output as follows: the first output is the Gröbner basis G of J_F , the second output is the list of leading monomials of elements in G , the third output is $M \setminus L(J_F)$ whose cardinality is precisely the fourth output. Since the problem is symmetric w.r.t. the y -axis, $\mu(F) = 2$ implies that each critical point has Milnor number 1.

Observe that the origin is not a critical point of F both as a polynomial map from $\mathbb{C}^3$ and from $\mathbb{C}^2$. In fact

```
> Milnor(F);
[[-(x-1)(x-2) - x(x-2) - x(x-1), 2y], [1, y], [], 0]
> Milnor(F, {x, y, z})
[[0, -(x-1)(x-2) - x(x-2) - x(x-1), 2y], [1, 1, y], [], 0]
```

In this case the first output is a standard bases of J_F whose leading monomials, w.r.t. a fixed l.m.o., give the second output.

At last let us observe that the zero locus $F^{-1}(0)$ is smooth both as a subset of $\mathbb{C}^2$ and of $\mathbb{C}^3$, in fact

```
> PolyTyurina(F);
[[1], [1], [], 0]
> PolyTyurina(F, {x, y, z});
[[1], [1], [], 0]
```

5.3. Optional input: some more subtle utilities. All the procedure described in 5.1.2 require three input: the polynomial F and two further optional input, precisely

- a set of variables, by default set as the variables `indets(F)` appearing in F ,
- a monomial order, by default set either as `tdeg(variables)`, which is the MAPLE command for the graduated reverse lexicographic g.m.o., or as `tdeg_min(variables)`, which is the l.m.o. defined as the `tdeg` opposite: the former is clearly introduced in `PolyMilnor` and `PolyTyurina` and the latter in `Milnor` and `Tyurina`.

Introducing different choices may show interesting possibilities of our subroutine.

5.3.1. Monomial ordering. It is a well known fact that the graduated reverse lexicographic g.m.o. is in general the more efficient monomial ordering for Buchberger algorithm: this is the reason for the default choices in `PolyMilnor` and `PolyTyurina`. Anyway, if needed, these procedures may run with many further g.m.o.: e.g. if, for any reason, the user will prefer to run `PolyTyurina` w.r.t. the pure lexicographic g.m.o. he will have to type, in the following case of a deformation of a threefold E_8 singularity:

```

> F:=x^2+y^3+z^5+t^2+y*z+y^2+z^2+y*z^2+z^3+y*z^3+z^4;
      F := x^2 + y^3 + z^5 + t^2 + yz^2 + z^3 + yz^3 + z^4
> PolyMilnor(F);
      [[x, t, 3yz^2 - 15y^2z + 4z^2 + 2yz + 3y^2, 3y^2 + z^2 + z^3,
      3375y^4 - 1677y^2z - 639y^3 + 224z^2 + 124yz + 114y^2,
      225y^3z - 324y^2z - 18y^3 + 88z^2 + 38yz + 93y^2],
      [x, t, yz^2, z^3, y^4, y^3z], [1, y, z, y^2, yz, z^2, y^3, y^2z], 8]
> PolyMilnor(F,plex(x,y,z,t));
      [[t, 31z^3 + 88z^4 + 159z^5 + 129z^6 + 75z^7,
      128z^3 + 319z^4 + 237z^5 + 225z^6 + 69z^2 + 46yz, 3y^2 + z^2 + z^3, x],
      [t, z^7, yz, y^2, x], [1, z, z^2, z^3, z^4, z^5, z^6, y], 8]

```

Observe how different are the two Gröbner bases and consequently the leading monomial bases and associated bases of quotient vector spaces. The user may also verify how much slower is `plex` w.r.t. the default `tdeg` by running by himself the routines.

In particular, running `Tyurina` w.r.t. different l.m.o.'s gives *different monomial basis of the Kuranishi space*:

```

> TyurinaBasis(F);
      [z^2, z, y, 1]
> TyurinaBasis(F,plex_min(x,y,z,t));
      [y^2, y, z, 1]

```

Remark 5.5. For what concerns efficiency of l.m.o.'s in `Milnor` and `Tyurina` it turns out that sometimes `plex_min` is more efficient than `tdeg_min`, as observed in the last Section 6 when proving Theorems 6.10 and 6.11. But we do not know if this is a general fact, then we keep `tdeg_min` as the default l.m.o. both in `Milnor` and `Tyurina`, for coherence with the default choice of `tdeg` for their global counterparts.

5.3.2. Variables. The default choice for the optional input `variables` as the set of those variables appearing in the given polynomial F (`indets(F)`) has been thought to make our routine more user friendly. Anyway this choice may hide some important subtleties, as already pointed out in Remark 5.4 in the case `variables` has been chosen as a greater set of variables w.r.t. `indets(F)`. Here we want to underline a significant potentiality of our routine when `variables` is chosen to be a *strictly smaller subset of indets(F)*.

Let us set $F = x^3 + x^4 + xy^2$. Then we get:

```

> F :=x^3+y^4+x*y^2;
> Milnor(F);
      [{6x^3 - 4y^4, 3x^2 + y^2, 4y^3 + 2xy}, [y^2, xy, x^3], [x^2, x, y, 1], 4]
> Tyurina(F);
      [{x^3 + y^4 + xy^2, 3x^2 + y^2, 4y^3 + 2xy, 2x^3 - y^4}, [y^2, xy, xy^2, x^3], [x^2, x, y, 1], 4]

```

Then $T^1 \cong \langle x^2, x, y, 1 \rangle_{\mathbb{C}}$ and

$$(21) \quad F_t := F + tx^2 = x^3 + y^4 + xy^2 + tx^2$$

is a non-trivial 1-parameter small deformation of F such that, for any fixed $t \in \mathbb{C}$, F_t has a critical point in $0 \in \mathbb{C}^2$ which is also a singular point of the plane curve $F_t^{-1}(0)$. We are interested in studying Milnor and Tyurina numbers of this singularity *for any* $t \in \mathbb{C}$. This can be performed by a careful use of the variables input. Let us first of all observe that if no optional input are added then we get

```
> Ft := F+t*x^2;
```

$$Ft := x^3 + y^4 + xy^2 + tx^2$$

```
> Milnor(Ft);
```

Error, (in Milnor) the given critical point is not isolated

In fact, by default `Milnor` considers F_t as a polynomial map defined over $\mathbb{C}^3(x, y, t)$. By forcing `Milnor` to work with variables $\{x, y\}$ only, then F_t is considered as a polynomial map defined over $\mathbb{C}^2$ with coefficient ring $\mathbb{C}[t]$, i.e. $F_t \in (\mathbb{C}[t])[x, y]$, giving:

```
> Milnor(Ft, {x, y})
```

$$[\{3x^2 + y^2 + 2tx, (1-4t)y^3 + 3yx^2, 4y^3 + 2xy\}, [x, y^3, xy], [y^2, y, 1], 3]$$

```
> Tyurina(Ft, {x, y})
```

$$[\{3x^2 + y^2 + 2tx, x^3 + y^4 + xy^2 + tx^2, (1-4t)y^3 + 3yx^2, 4y^3 + 2xy\}, [x, y^3, x^2, xy], [y^2, y, 1], 3]$$

This means that, *for generic*¹ t , $\tau_{F_t}(0) = 3 = \mu_{F_t}(0)$. Moreover *by looking at the leading coefficients of the given standard basis of J_F we get all the relations defining non-generic values for t* , precisely:

```
> MB := MilnorGroebnerBasis(Ft, {x, y})
```

$$MB := \{3x^2 + y^2 + 2tx, (1-4t)y^3 + 3yx^2, 4y^3 + 2xy\}$$

```
> for i from 1 to nops(MB) do
```

```
> LeadingTerm(MB[i], tdeg_min(x,y)) end do;
```

$$2t, x$$

$$1-4t, y^3$$

$$2, xy$$

For $t = 0$ we do not have any deformation of F , giving $\tau_{F_0}(0) = 4 = \mu_{F_0}$. But the further relation $1-4t = 0$ gives:

```
> t := 1/4: Milnor(Ft, {x, y})
```

$$[\{3x^2 + y^2 + 1/2x, 36y^3x^2 + 12y^5, 4y^3 + 2xy\}, [x, y^5, xy], [y^4, y^3, y^2, y, 1], 5]$$

```
> Tyurina(Ft, {x, y})
```

$$\left[\left\{ x^3 + y^4 + xy^2 + 1/4x^2, \frac{3}{64}y^4x^2 + \frac{1}{64}y^6 - \frac{27}{256}x^6 - \frac{9}{256}x^4y^2, 4y^3 + 2xy, 36y^3x^2 + 12y^5, 3x^2 + y^2 + 1/2x \right\}, [x, y^6, x^2, y^5, xy], [y^4, y^3, y^2, y, 1], 5 \right]$$

Therefore

$$\tau_{F_t}(0) = \mu_{F_t}(0) = \begin{cases} 5 & \text{for } t = 1/4, \\ 4 & \text{for } t = 0, \\ 3 & \text{otherwise.} \end{cases}$$

¹ t is treated as a variable without any evaluation.

In particular Proposition 3.9 implies that, *for any* t , $0 \in \text{Spec } \mathcal{O}_{F_t,0}$ is a *w.h. singularity*, in spite of the fact that F_t is never a w.h. polynomial.

5.4. An efficiency remark: global to local subroutines. After numerous applications of the previous routines the reader will convince himself that the **SB** procedure turns out to be less efficient than the Buchberger algorithm as implemented in MAPLE. As a consequence our routines can be arranged in the following decreasing sequence of efficiency:

$$\text{PolyMilnor} > \text{PolyTyurina} > \text{Milnor} > \text{Tyurina} .$$

A slight improvement of **Milnor** and **Tyurina** efficiency can be obtained by applying the **SB** algorithm to a Gröbner basis of J_f and I_f rather than to their original generators. What is obtained is a sort of “pasting” of global and local routines, precisely:

- **MILNOR** which is a procedure computing Milnor numbers of both a polynomial f and of a critical point of f :

Input : $F \in \mathbb{C}[\mathbf{x}]$, **Basis** in the **Groebner** package , **SB** ,
Optional: **variables**, by default assigned by **indets(F)**
Optional: a couple [l.m.o., g.m.o.], by default assigned by
[**tdeg_min(variables)**, **tdeg(variables)**]

Output : a Gröbner basis G of $J_F \subset (\mathbb{C}[\mathbf{x} \setminus \text{variables}])[\text{variables}]$, $L_{Gl}(G)$,
 $M \setminus L_{Gl}(J_F)$, $\mu(F)$,
a standard basis B of $J_F \subset (\mathbb{C}[\mathbf{x} \setminus \text{variables}])[\text{variables}]$, $L_{Lo}(B)$,
 $M \setminus L_{Lo}(J_F)$, $\mu_F(0)$;

- **TYURINA** which is a the procedure computing the Tyurina numbers of both a polynomial f and of a critical point of f :

Input : $F \in \mathbb{C}[\mathbf{x}]$, **Basis** in the **Groebner** package , **SB** ,
Optional: **variables**, by default assigned by **indets(F)**
Optional: a couple [l.m.o., g.m.o.], by default assigned by
[**tdeg_min(variables)**, **tdeg(variables)**]

Output : a Gröbner basis G of $J_F \subset (\mathbb{C}[\mathbf{x} \setminus \text{variables}])[\text{variables}]$, $L_{Gl}(G)$,
 $M \setminus L_{Gl}(I_F)$, $\tau(F)$,
a standard basis B of $I_F \subset (\mathbb{C}[\mathbf{x} \setminus \text{variables}])[\text{variables}]$, $L_{Lo}(B)$,
 $M \setminus L_{Lo}(I_F)$, $\tau_F(0)$;

where L_{Lo} and L_{Gl} mean *leading monomials w.r.t the given local and global m.o.*, respectively.

5.4.1. The MAPLE details. Let us start with the procedure computing Milnor numbers:

The associated sub-procedures:

```
> PolyMILNORGroebnerBasis := proc (F,variables::set:=indets(F),
> SMOS::anything:=[tdeg_min(op(variables)),tdeg(op(variables))])
> MILNOR(F,variables,SMOS)[1] end proc:
```

```

> PolyMILNORLT := proc (F, variables::set:= indets(F),
> SMOS::anything:=[tdeg_min(op(variables)),tdeg(op(variables))])
> MILNOR(F, variables, SMOS)[2] end proc:

> PolyMILNORBasis := proc (F,variables::set := indets(F),
> SMOS::anything:=[tdeg_min(op(variables)),tdeg(op(variables))])
> MILNOR(F,variables,SMOS)[3] end proc:

> PolyMILNORNumber:=proc (F, variables::set:= indets(F),
> SMOS::anything:=[tdeg_min(op(variables)),tdeg(op(variables))])
> MILNOR(F,variables,SMOS)[4] end proc:

> MILNORGroebnerBasis := proc(F,variables::set := indets(F),
> SMOS::anything:=[tdeg_min(op(variables)),tdeg(op(variables))])
> MILNOR(F,variables,SMOS)[5] end proc:

> MILNORLT := proc (F, variables::set := indets(F),
> SMOS::anything:=[tdeg_min(op(variables)),tdeg(op(variables))])
> MILNOR(F, variables, SMOS)[6] end proc:

> MILNORBasis := proc (F,variables::set :=indets(F),
> SMOS::anything:=[tdeg_min(op(variables)),tdeg(op(variables))])
> MILNOR(F, variables,SMOS)[7] end proc:

> MILNORNumber:=proc (F,variables::set :=indets(F),
> SMOS::anything:=[tdeg_min(op(variables)),tdeg(op(variables))])
> MILNOR(F,variables,SMOS)[8] end proc:

```

Then the routine computing Tyurina numbers:

```

> TYURINA := proc (F, variables::set := indets(F),
> SMOS::anything := [tdeg_min(op(variables)),tdeg(op(variables))])
> U := SMOS[1]; V := SMOS[2];
> if type(U, ShortMonomialOrder) = false then error
> "invalid input: the first ShortMonomialOrder is not well defined"
> else if localorglobal(U, variables) <> Lo then error
> "invalid input: the first ShortMonomialOrder should be LOCAL"
> else if type(V, ShortMonomialOrder) = false then error
> "invalid input: the second ShortMonomialOrder is not well defined"
> else if localorglobal(V, variables) <> Gl then error
> "invalid input: the second ShortMonomialOrder should be GLOBAL"
> else r := nops(variables);
> J := [F, seq(diff(F, variables[i]), i = 1 .. r)];
> J := Basis(J, V);
> IniJ := [seq(LeadingMonomial(J[i], V), i = 1 .. nops(J))];
> massimo:=r*max(seq(degree(IniJ[i],variables),i = 1..nops(IniJ)));
> if IsZeroDimensional(<(op(IniJ))>,{op(variables)})= false then error
> "the given singular point is not isolated"
> else LUU := [op(subs(vq = 1, convert(map(expand,
> series(1/(product(1-variables[k]*vq, k = 1 .. r)),
> vq, massimo+1)), polynom))];
> L := sort(LUU, proc (t1, t2) options operator, arrow;
> TestOrder(t1, t2, V) end proc);
> N := [];
> for j to nops(L) do up := 1;
> for k to nops(IniJ) do
> if divide(L[j], IniJ[k]) then up := 0
> end if end do;
> if up = 1 then N := [op(N),L[j]]
> end if end do;

```

```

> G := SB(J, variables, U);
> Ini := [seq(LeadingMonomial(G[i], U), i = 1 .. nops(G))];
> massimo := r*max(seq(degree(Ini[i], variables), i = 1 .. nops(Ini)));
> LUU := [op(subs(vq = 1, convert(map(expand,
> series(1/(product(1-variables[h]*vq, h = 1 .. r)),
> vq, massimo+1)), polynom))];
> L := sort(LUU, proc (t1, t2) options operator, arrow;
> TestOrder(t1, t2, U) end proc);
> M := [];
> for j to nops(L) do up := 1;
> for k to nops(Ini) do
> if divide(L[j], Ini[k]) then up :=
> 0 end if end do;
> if up = 1 then M := [op(M), L[j]]
> end if end do;
> [J, IniJ, N, nops(N), G, Ini, M, nops(M)]
> end if end if end if end if end if end if end proc:

```

The associated sub-procedures:

```

> PolyTYURINAGroebnerBasis := proc(F, variables::set := indets(F),
> SMOS::anything := [tdeg_min(op(variables)), tdeg(op(variables))])
> TYURINA(F, variables, SMOS)[1] end proc:

> PolyTYURINALT := proc (F, variables::set := indets(F),
> SMOS::anything := [tdeg_min(op(variables)), tdeg(op(variables))])
> TYURINA(F, variables, SMOS)[2] end proc:

> PolyTYURINABasis := proc (F, variables::set := indets(F),
> SMOS::anything := [tdeg_min(op(variables)), tdeg(op(variables))])
> TYURINA(F, variables, SMOS)[3] end proc:

> PolyTYURINANumber := proc (F, variables::set := indets(F),
> SMOS::anything := [tdeg_min(op(variables)), tdeg(op(variables))])
> TYURINA(F, variables, SMOS)[4] end proc:

> TYURINAGroebnerBasis := proc(F, variables::set := indets(F),
> SMOS::anything := [tdeg_min(op(variables)), tdeg(op(variables))])
> TYURINA(F, variables, SMOS)[5] end proc:

> TYURINALT := proc (F, variables::set := indets(F),
> SMOS::anything := [tdeg_min(op(variables)), tdeg(op(variables))])
> TYURINA(F, variables, SMOS)[6] end proc:

> TYURINABasis := proc (F, variables::set := indets(F),
> SMOS::anything := [tdeg_min(op(variables)), tdeg(op(variables))])
> TYURINA(F, variables, SMOS)[7] end proc:

> TYURINANumber := proc (F, variables::set := indets(F),
> SMOS::anything := [tdeg_min(op(variables)), tdeg(op(variables))])
> TYURINANor(F, variables, SMOS)[8] end proc:

```

5.4.2. *An example of application.* Let us consider the same polynomial $F(x, y) = x^3 + y^4 + xy^2$ given in 5.3.2. Procedures MILNOR and TYURINA give *all* the information we could get by applying all the introduced routines, precisely:

```

> MILNOR(F)
[[3 x^2 + y^2, 2 y^3 + xy], [x^2, y^3], [1, y, x, y^2, xy, xy^2], 6,
{-2 y^4 + 3 x^3, 2 y^3 + xy, 3 x^2 + y^2}, [y^2, xy, x^3], [x^2, x, y, 1], 4]
> TYURINA(F)

```

$$\begin{aligned} & [[xy, 3x^2 + y^2, y^3], [xy, x^2, y^3], [1, y, x, y^2], 4, \\ & \{xy, 3x^3, 3x^2 + y^2, y^3\}, [y^2, x^3y^3, xy], [x^2, x, y, 1], 4] \end{aligned}$$

In particular it turns out that F admits some further critical point which is not a singular point of $F^{-1}(0)$. By

```
> solve({diff(F,x),diff(F,y)},{x,y});
```

$$[[x = 0, y = 0], [x = 0, y = 0], [x = -1/2 \left(\text{RootOf}(3_Z^2 + 1, \text{label} = _L1) \right)^2, \\ y = 1/2 \text{RootOf}(3_Z^2 + 1, \text{label} = _L1)], [x = 0, y = 0]]$$

it follows that there are precisely two further critical points of F having Milnor number 1 and Tyurina number 0. Let us now type:

```
> TyB := TyurinaBasis(F)
      TyB := [x^2, x, y, 1]
> T := nops(TyB):
> F[Lambda] := F+sum(lambda[i]*TyB[T-i],i= 0 .. T-1);
      FΛ := x3 + y4 + xy2 + λ0 + λ1y + λ2x + λ3x2
> TYURINA(F[Lambda], {x, y});
      [[1], [1], [], 0, [1], [1], [], 0]
> Tyurina(F[Lambda], {x, y});
Warning, computation interrupted
```


We had to interrupt the calculation of `Tyurina` since it wasn't able to produce any output after considerable time, on the contrary of `TYURINA` which quickly produced the given (trivial) output.

Remark 5.6. What observed in 5.2 and 5.3, about optional inputs for `Milnor` and `Tyurina`, applies analogously for `MILNOR` and `TYURINA`.

6. APPLICATION: ADJACENCIES OF ARNOL'D SIMPLE SINGULARITIES

Let us consider the classes of Arnol'd simple singularities A_n, D_n, E_6, E_7, E_8 . Recall that a class of singularities B is said to be *adjacent* to a class of singularities A (notation $A \leftarrow B$) if any singularity in B can be deformed to a singularity in A by an arbitrarily small deformation (see [4] §I.2.7, [5] §15.0 and [16] §7.C). Adjacency turns out to be a *partial order relation* on the set of singularities' equivalence classes. In the following we will employ the optional input on variables, as observed in 5.3.2, to show explicit equations of algebraic *stratifications* of Kuranishi spaces verifying the following Arnol'd adjacency diagram

(22)



(see [3], [5] and in particular [4] §I.2.7). Such a stratification gives a geometric interpretation, by means of inclusions of algebraic subsets, of the partial order relation induced by adjacency.

Let us first of all observe that the Kuranishi space T^1 of an Arnol'd simple m -fold singularity of type A_n, D_n, E_6, E_7, E_8 do not depend on its dimension m , since partials of quadratic terms are linear generators of I_f eliminating the associated variable from $M \setminus L(I_f)$. Therefore the study of diagram (22) can be reduced to the case of Arnol'd simple *curve* singularities whose local equations are given in (17): this is actually guaranteed by the following Morse Splitting Lemma 6.2.

In the sequel we will need the following notation and results, essentially due to V.I. Arnol'd [5]. We refer the interested reader to books [4], [5], [16] and [12] for details and proofs.

Definition 6.1 (Co-rank of a critical point). The *co-rank* of a critical point p of a holomorphic function f defined over an open subset of $\mathbb{C}^n$ is the number

$$\text{crk}_f(p) := n - \text{rk}(\text{Hess}_f(p))$$

where $\text{Hess}_f(p)$ is the Hessian matrix of f in p (for shortness the function f will be omitted when clear from the context). In particular if $\text{crk}(p) = 0$ then p is called a *non-degenerate*, or *Morse*, *critical point*.

Lemma 6.2 (Morse Splitting Lemma, [5] §11.1, [16] (7.16) and [12] Theorem I.2.47). Assume that $f \in \mathfrak{m}^2 \subset \mathbb{C}\{\mathbf{x}\}$ and $\text{crk}_f(0) = n - k$. Then f is equivalent² to the following germ of singularity

$$\sum_{i=1}^k x_i^2 + g(x_{k+1}, \dots, x_n)$$

where $g \in \mathfrak{m}^3$ is uniquely determined (up to equivalence).

Theorem 6.3 ([12], Theorems I.2.46, I.2.48, I.2.51, I.2.53).

- (1) For $f \in \mathfrak{m}^2 \subset \mathbb{C}\{\mathbf{x}\}$ the following facts are equivalent:
 - $\text{crk}(0) = 0$ i.e. 0 is a non-degenerate critical point of f ,
 - $\mu(0) = 1 = \tau(1)$,
 - $0 \in f^{-1}(0)$ is, up to equivalence, a node i.e. a simple A_1 singularity.
- (2) For $f \in \mathfrak{m}^2 \subset \mathbb{C}\{\mathbf{x}\}$ the following facts are equivalent:
 - $\text{crk}(0) \leq 1$ and $\mu(0) = m$,
 - $0 \in f^{-1}(0)$ is equivalent to a simple A_m singularity.
- (3) For $f \in \mathfrak{m}^3 \subset \mathbb{C}\{y, z\}$ the following facts are equivalent:
 - the 3-jet $f^{(3)}$ of f factors into at least two distinct linear factors and $\mu(0) = m \geq 4$,
 - $0 \in f^{-1}(0)$ is equivalent to a simple D_m singularity.
- (4) For $f \in \mathfrak{m}^3 \subset \mathbb{C}\{y, z\}$ the following facts are equivalent:
 - the 3-jet $f^{(3)}$ of f has a unique liner factor (of multiplicity 3) and $\mu(f) \leq 8$,
 - $0 \in f^{-1}(0)$ is equivalent to a simple singularity of type E_6, E_7 or E_8 and $\mu(0) = 6, 7$ or 8 respectively.

²In the sense of Definition 3.7 and Remark 3.8.

Let us now introduce a non-standard notation, useful to describe a nice geometric property of stratifications via algebraic subsets of a simple Arnol'd singularity's Kuranishi space, as explained in the following statements. Consider the following square of subset inclusions

$$(23) \quad \begin{array}{ccc} B & \hookrightarrow & D \\ \uparrow & & \uparrow \\ A & \hookrightarrow & C \end{array}$$

Then $A \subseteq B \cap C$, necessarily.




FIGURE 1. The c.i.p. for the inclusions' square (23)

Definition 6.4 (Complete Intersection Property - (c.i.p.)). A square of subset inclusions (23) is said to admit the *complete intersection property* if

$$A = B \cap C.$$

For shortness we will say that (23) is a *c.i.p. square*. The geometric meaning of c.i.p. in (23) is explained by Figure 1, while Figure 2 describes geometrically the following *sequence of two c.i.p. squares*

$$(24) \quad \begin{array}{ccccc} B & \hookrightarrow & D & \hookrightarrow & F \\ \uparrow & & \uparrow & & \uparrow \\ A & \hookrightarrow & C & \hookrightarrow & E \end{array}$$

meaning that $A = B \cap C = B \cap D \cap E$. Moreover the following inclusions' diagram

$$(25) \quad \begin{array}{ccccc} B \cup E & \xleftarrow{\quad} & E & & \\ \uparrow & \searrow & & \searrow & \\ A & & D & \xleftarrow{\quad} & G \\ & \searrow & \uparrow & & \\ & & C & & \end{array}$$




FIGURE 2. The sequence (24) of two c.i.p. squares

is called a *union of c.i.p. squares* if $A = B \cap C$ and $E = F \cap G$. A particular case, occurring in the following, is when $C = G$: then diagram (25) becomes the following one

$$(26) \quad \begin{array}{ccc} B \cup F & \xrightarrow{\quad} & D \\ \uparrow & & \uparrow \\ E & \xrightarrow{\quad} & C = G \\ \cup & \nearrow & \\ A & & \end{array}$$

and we will say this diagram to represent a *hinged union of c.i.p. squares*, whose hinge is the inclusion $C \hookrightarrow D$.

At last the following inclusions' diagram

$$(27) \quad \begin{array}{ccc} B & \xrightarrow{\quad} & D \\ \uparrow & & \uparrow \\ E & \xrightarrow{\quad} & C \\ \cup & \nearrow & \\ A & & \end{array}$$

is called a *reducible c.i.p. square* if $A \cup E = B \cap C$ i.e. if

$$\begin{array}{ccc} B & \xrightarrow{\quad} & D \\ \uparrow & & \uparrow \\ A \cup E & \xrightarrow{\quad} & C \end{array}$$

is a c.i.p. square.

6.1. Outline of the following results. Statements and proofs of the following Theorems 6.5 , 6.6 , 6.8 , 6.10 and 6.11 have the same structure we are going to outline here. Precisely their statements describe set theoretical stratifications by algebraic subsets of the Kuranishi space T^1 of simple hypersurface singularities

A_n, D_n and E_n (the latter with $6 \leq n \leq 8$). Their proofs go on by the following steps:

- (1) look for the critical points of a generic small deformation of our initial simple singularity, by solving the polynomial system assigned by partial derivatives (Jacobian ideal generators): they turn out to be precisely n points (with $6 \leq n \leq 8$ in the E_n case);
- (2) imposing one of the previous critical points to be actually a singular point means defining a hypersurface $\mathcal{L} \subset T^1$: we have then n of such hypersurfaces, one for each critical point;
- (3) any of those hypersurfaces is then stratified by nested algebraic subsets defined by a progressive vanishing of leading coefficients in Jacobian ideals' standard bases of more and more specialized deformations.

The last step (3) is obtained by a systematic use of routines **Milnor**, **Tyurina**, **MILNOR** and **TYURINA** previously described, running with suitable l.m.o.'s defined on a strict subset of variables appearing in the polynomial equation of a generic small deformation, as explained in 5.3.2. Such a procedure allows to explicitly write down relations on deformation parameters and then equations of the algebraic stratifications.

Caveat Lector. The large use of MAPLE routines in step (3) makes difficult to follow and understand algebraic arguments without the support of MAPLE 11 or 12. Hence the reader is warmly advised of repeating by himself MAPLE commands as described in the following. Moreover since the following results are here reported as a significant application of the previously presented subroutines, their role, inputs and outputs will be explained and listed in detail; on the contrary the use of standard MAPLE procedures, like e.g. **eliminate** or **EliminationIdeal**, will be left to the reader who may helpfully consult the excellent on-line MAPLE help system.

6.2. Simple singularities of A_n type.

Theorem 6.5. *Let T^1 be the Kuranishi space of a simple N -dimensional singular point $0 \in f^{-1}(0)$ with*

$$f(x_1, \dots, x_{N+1}) = \sum_{i=1}^N x_i^2 + x_{N+1}^{n+1} \quad (\text{for } n \geq 1) .$$

The subset of T^1 parameterizing small deformations of $0 \in f^{-1}(0)$ to a simple node (i.e. an A_1 singularity) is the union of n hypersurfaces. Moreover, calling $\mathcal{L}$ any of those hypersurfaces, there exists a stratification of nested algebraic subsets

$$(28) \quad \mathcal{L} \longleftarrow \supset \mathcal{V}_2 \longleftarrow \supset \dots \longleftarrow \supset \mathcal{V}_2^m \longleftarrow \supset \dots \longleftarrow \supset \mathcal{V}_2^n$$

verifying the Arnol'd's adjacency diagram

$$(29) \quad A_1 \longleftarrow A_2 \longleftarrow \dots \longleftarrow A_m \longleftarrow \dots \longleftarrow A_n$$

where

- $\mathcal{L}$ is the hypersurface of T^1 defined by equation (31), keeping in mind (30),
- $\mathcal{V}_2^m := \bigcap_{k=2}^m \mathcal{V}_k$ where $\mathcal{V}_k$ are hypersurfaces of $\mathcal{L}$ defined by the vanishing of variables v_k introduced by (33).

Proof. By the Morse Splitting Lemma 6.2 we can reduce to the case $N = 1$ with $f(y, z) = y^2 + z^{n+1}$ for $n \geq 1$. Then Proposition 2.3 gives

$$T^1 \cong \mathbb{C}[y, z]/(y, z^n) \cong \langle 1, z, \dots, z^{n-1} \rangle_{\mathbb{C}}$$

and, given $\Lambda = (\lambda_0, \dots, \lambda_{n-1}) \in T^1$, the associated deformation of $U_0 = \text{Spec}(\mathcal{O}_{f,0})$ is

$$U_{\Lambda} = \{f_{\Lambda}(y, z) := f(y, z) + \sum_{i=0}^{n-1} \lambda_i z^i = 0\}.$$

A solution of the jacobian system of partial derivatives is then given by $p_{\Lambda} = (0, z_{\Lambda})$ where z_{Λ} is a zero of the following polynomial

$$(30) \quad (n+1)z^n + \sum_{i=1}^{n-1} i\lambda_i z^{i-1} \in \mathbb{C}[\lambda][z].$$

This means that f_{Λ} admits precisely n critical points. Observe that imposing $p_{\Lambda} \in U_{\Lambda}$, which is asking for one of the previous critical points to be actually a singular point of U_{Λ} , defines the following hypersurface in T^1

$$(31) \quad p_{\Lambda} \in U_{\Lambda} \iff \Lambda \in \mathcal{L} := \{z_{\Lambda}^{n+1} + \sum_{i=0}^{n-1} \lambda_i z_{\Lambda}^i = 0\} \subset T^1.$$

Notice that we get precisely n such hypersurfaces, one for each critical point of f_{Λ} .

- $\mathcal{L}$ is a hypersurface of the Kuranishi space parameterizing small deformations of U_0 admitting a singular point of type at least A_1 in the origin.

After translating $z \mapsto z + z_{\Lambda}$, we get

$$(32) \quad \begin{aligned} f_{\Lambda}(y, z + z_{\Lambda}) = f(y, z) &+ \left(z_{\Lambda}^{n+1} + \sum_{i=0}^{n-1} \lambda_i z_{\Lambda}^i \right) \\ &+ \left((n+1)z_{\Lambda}^n + \sum_{i=1}^{n-1} i\lambda_i z_{\Lambda}^{i-1} \right) z \\ &+ \sum_{k=2}^{n-1} \left(\binom{n+1}{k} z_{\Lambda}^{n+1-k} + \sum_{i=k}^{n-1} \binom{i}{k} \lambda_i z_{\Lambda}^{i-k} \right) z^k \\ &+ (n+1)z_{\Lambda} z^n. \end{aligned}$$

Then by (31) and (30) the origin (i.e. $p_{\Lambda} \in U_{\Lambda}$) is at least a node: in fact by running Milnor and Tyurina e.g. when $n = 7$, we get (set $L := z_{\Lambda}$):

> F:=y^2 + z^8;

$$F := y^2 + z^8$$

> TyB := TyurinaBasis(F)

$$TyB := [z^6, z^5, z^4, z^3, z^2, z, 1]$$

> T := nops(TyB):

> F[Lambda] := F+sum(lambda[i]*TyB[T-i], i = 0 .. T-1)

$$F_{\Lambda} := y^2 + z^8 + \lambda_0 + \lambda_1 z + \lambda_2 z^2 + \lambda_3 z^3 + \lambda_4 z^4 + \lambda_5 z^5 + \lambda_6 z^6$$

> solve({diff(F[Lambda], y), diff(F[Lambda], z)}, [y, z])

$$[[y = 0, z = \text{RootOf}(8_Z^7 + \lambda_1 + 2\lambda_2_Z + 3\lambda_3_Z^2 + 4\lambda_4_Z^3 + 5\lambda_5_Z^4 + 6\lambda_6_Z^5)]]$$

Notice that the polynomial in *RootOf* is precisely (30) for $n = 7$.

```
> z := Z+L:
> collect(F[Lambda], [y, Z], 'distributed')

$$y^2 + Z^8 + 8LZ^7 + (\lambda_6 + 28L^2)Z^6 + (56L^3 + \lambda_5 + 6\lambda_6L)Z^5 +$$


$$(70L^4 + 5\lambda_5L + 15\lambda_6L^2 + \lambda_4)Z^4 + (10\lambda_5L^2 + \lambda_3 + 56L^5 + 4\lambda_4L + 20\lambda_6L^3)Z^3$$


$$+ (10\lambda_5L^3 + 3\lambda_3L + 6\lambda_4L^2 + 15\lambda_6L^4 + \lambda_2 + 28L^6)Z^2 +$$


$$(4\lambda_4L^3 + 8L^7 + 5\lambda_5L^4 + 6\lambda_6L^5 + 2\lambda_2L + 3\lambda_3L^2 + \lambda_1)Z$$


$$+ \lambda_2L^2 + \lambda_5L^5 + \lambda_6L^6 + L^8 + \lambda_0 + \lambda_3L^3 + \lambda_4L^4 + \lambda_1L$$

```

Compare (32) for $n=7$ with the previous output expression.

```
> SD := [seq(Z^i, i = 2 .. 7)]
SD := [Z^2, Z^3, Z^4, Z^5, Z^6, Z^7]
> for j from 2 to 7 do
> FL[j] := y^2 + Z^8 + sum(v[i]*SD[i-1], i = j .. 7) end do
FL2 := y^2 + Z^8 + v2Z^2 + v3Z^3 + v4Z^4 + v5Z^5 + v6Z^6 + v7Z^7
FL3 := y^2 + Z^8 + v3Z^3 + v4Z^4 + v5Z^5 + v6Z^6 + v7Z^7
FL4 := y^2 + Z^8 + v4Z^4 + v5Z^5 + v6Z^6 + v7Z^7
FL5 := y^2 + Z^8 + v5Z^5 + v6Z^6 + v7Z^7
FL6 := y^2 + Z^8 + v6Z^6 + v7Z^7
FL7 := y^2 + Z^8 + v7Z^7
> for j from 2 to 7 do Milnor(FL[9-j], {y, Z}) end do
[[2y, 8Z^7 + 7v7Z^6], [y, Z^6], [Z^5, Z^4, Z^3, Z^2, Z, 1], 6]
[[2y, 8Z^7 + 6v6Z^5 + 7v7Z^6], [y, Z^5], [Z^4, Z^3, Z^2, Z, 1], 5]
[[2y, 8Z^7 + 5v5Z^4 + 6v6Z^5 + 7v7Z^6], [y, Z^4], [Z^3, Z^2, Z, 1], 4]
[[2y, 8Z^7 + 4v4Z^3 + 5v5Z^4 + 6v6Z^5 + 7v7Z^6], [y, Z^3], [Z^2, Z, 1], 3]
[[2y, 8Z^7 + 3v3Z^2 + 4v4Z^3 + 5v5Z^4 + 6v6Z^5 + 7v7Z^6], [y, Z^2], [Z, 1], 2]
[[2y, 8Z^7 + 2v2Z + 3v3Z^2 + 4v4Z^3 + 5v5Z^4 + 6v6Z^5 + 7v7Z^6], [y, Z], [1], 1]
> for j from 2 to 7 do Tyurina(FL[9-j], {y, Z}) end do
[[y^2 + Z^8 + v7Z^7, 2y, 8Z^7 + 7v7Z^6], [y^2, y, Z^6], [Z^5, Z^4, Z^3, Z^2, Z, 1], 6]
[[y^2 + Z^8 + v6Z^6 + v7Z^7, 2y, 8Z^7 + 6v6Z^5 + 7v7Z^6], [y^2, y, Z^5], [Z^4, Z^3, Z^2, Z, 1], 5]
[[y^2 + Z^8 + v5Z^5 + v6Z^6 + v7Z^7, 2y, 8Z^7 + 5v5Z^4 + 6v6Z^5 + 7v7Z^6], [y^2, y, Z^4],
[Z^3, Z^2, Z, 1], 4]
[[y^2 + Z^8 + v4Z^4 + v5Z^5 + v6Z^6 + v7Z^7, 2y, 8Z^7 + 4v4Z^3 + 5v5Z^4 + 6v6Z^5 + 7v7Z^6],
[y^2, y, Z^3], [Z^2, Z, 1], 3]
[[y^2 + Z^8 + v3Z^3 + v4Z^4 + v5Z^5 + v6Z^6 + v7Z^7, 2y, 8Z^7 + 3v3Z^2 + 4v4Z^3 + 5v5Z^4 +
6v6Z^5 + 7v7Z^6], [y^2, y, Z^2], [Z, 1], 2]
Warning, computation interrupted
> for j from 2 to 7 do TYURINA(FL[9-j], {y, Z}) end do
[[y, Z^6], [y, Z^6], [1, Z, Z^2, Z^3, Z^4, Z^5], 6, [y, Z^6], [y, Z^6], [Z^5, Z^4, Z^3, Z^2, Z, 1], 6]
```

$$\begin{aligned}
& [[y, Z^5], [y, Z^5], [1, Z, Z^2, Z^3, Z^4], 5, [y, Z^5], [y, Z^5], [Z^4, Z^3, Z^2, Z, 1], 5] \\
& [[y, Z^4], [y, Z^4], [1, Z, Z^2, Z^3], 4, [y, Z^4], [y, Z^4], [Z^3, Z^2, Z, 1], 4] \\
& [[y, Z^3], [y, Z^3], [1, Z, Z^2], 3, [y, Z^3], [y, Z^3], [Z^2, Z, 1], 3] \\
& [[y, Z^2], [y, Z^2], [1, Z], 2, [y, Z^2], [y, Z^2], [Z, 1], 2] \\
& [[Z, y], [Z, y], [1], 1, [Z, y], [Z, y], [1], 1]
\end{aligned}$$

We reported here both the output obtained with **Tyurina** and **TYURINA** to give a further account of differences between the two procedures: after some time, we had to stop **Tyurina** during the calculation of the last output, while **TYURINA** was able to quickly conclude the calculation; in fact the standard bases of I_{FL_j} obtained by the latter procedure are noticeably simpler than those obtained by the former procedure. In the following we will employ **TYURINA** every time **Tyurina** will be inefficient.

Fist of all let us observe that, for any $j = 2, \dots, 7$, $\tau(0) = \mu(0)$ meaning that the origin deforms always as a w.h. singularity, by Proposition 3.9. Moreover $\text{crk}(0) \leq 1$, since the rank of the Hessian matrix is always at least 1 for the contribution of y^2 . Then *the origin deforms always as a simple A_μ singularity, where μ is its Milnor number* by Theorem 6.3(2). In particular the standard basis of J_{FL_2} is given by $\{2y, 8Z^7 + 2v_2Z + 3v_3Z^2 + 4v_4Z^3 + 5v_5Z^4 + 6v_6Z^5 + 7v_7Z^6\}$ whose leading coefficient w.r.t. the default l.m.o. is $2v_2$. This means that

- $v_2 = 0$ is the equation of the codimension 1 subvariety $\mathcal{V}_2 \subset \mathcal{L}$ parameterizing small deformation of U_0 admitting a singularity of type at least A_2 in the origin.

Analogously the standard basis of J_{FL_3} is $\{2y, 8Z^7 + 3v_3Z^2 + 4v_4Z^3 + 5v_5Z^4 + 6v_6Z^5 + 7v_7Z^6\}$ whose leading coefficient is $3v_3$ meaning that

- $v_3 = 0$ is the equation of the subvariety $\mathcal{V}_3 \subset \mathcal{L} \subset T^1$ such that $\mathcal{V}_2 \cap \mathcal{V}_3 \subset \mathcal{L}$ is the codimension 2 subvariety parameterizing small deformation of U_0 admitting a singularity of type at least A_3 in the origin,

and so on: in general, for $n \geq 3$, setting

$$\begin{aligned}
(33) \quad v_k &:= \binom{n+1}{k} z_\Lambda^{n+1-k} + \sum_{i=k}^{n-1} \binom{i}{k} \lambda_i z_\Lambda^{i-k} \quad , \quad k = 2, \dots, n-1 \\
v_n &:= z_\Lambda
\end{aligned}$$

and defining codimension 1 subvarieties $\mathcal{V}_k := \{v_k = 0\}$ of $\mathcal{L}$, then

- $p_\Lambda \in U_\Lambda$ turns out to be a A_m ($2 \leq m \leq n$) simple hypersurface singularity if and only if Λ is the generic element of the codimension m subvariety $\mathcal{V}_2^m := \bigcap_{k=2}^m \mathcal{V}_k \subset T^1$.

This gives precisely the nested stratification (28) verifying the top row (29) in diagram (22) of Arnol'd's adjacencies. $\square$

6.3. Simple singularities of D_n type.

Theorem 6.6. *Let T^1 be the Kuranishi space of a simple N -dimensional singular point $0 \in f^{-1}(0)$ with*

$$f(x_1, \dots, x_{N+1}) = \sum_{i=1}^{N-1} x_i^2 + x_N^2 x_{N+1} + x_{N+1}^{n-1} \quad (\text{for } n \geq 4)$$

The subset of T^1 parameterizing small deformations of $0 \in f^{-1}(0)$ to a simple node is the union of n hypersurfaces. Moreover, calling $\mathcal{L}$ any of those hypersurfaces, there exists a stratification of nested algebraic subsets giving rise to the following sequence of inclusions and c.i.p. squares

$$(34) \quad \begin{array}{ccccccc} \mathcal{L} & \longleftarrow & \mathcal{W}_2 & \longleftarrow & \mathcal{V}_0^1 \cup \mathcal{W}_2^3 & \longleftarrow \dots \mathcal{W}_2^m \dots & \longleftarrow & \mathcal{W}_2^{n-1} \\ & & & & \uparrow & & \uparrow & \uparrow \\ & & & & \mathcal{V}_0^2 & \longleftarrow \dots \mathcal{V}_0^{m-1} \dots & \longleftarrow & \mathcal{V}_0^{n-2} = \{0\} \end{array}$$

verifying the Arnol'd's adjacency diagram

$$(35) \quad \begin{array}{ccccccc} A_1 & \longleftarrow & A_2 & \longleftarrow & A_3 & \longleftarrow \dots A_m \dots & \longleftarrow & A_{n-1} \\ & & & & \uparrow & & \uparrow & \uparrow \\ & & & & D_4 & \longleftarrow \dots D_m \dots & \longleftarrow & D_n \end{array}$$

where

- $\mathcal{L}$ is the hypersurface of T^1 defined by equation (38), keeping in mind (37),
- $\mathcal{V}_0^m := \bigcap_{k=0}^m \mathcal{V}_k$ and $\mathcal{W}_2^m := \bigcap_{k=2}^m \mathcal{W}_k$ where $\mathcal{V}_k, \mathcal{W}_k$ are hypersurfaces of $\mathcal{L}$ defined by equations (47), keeping in mind definitions (46).

Remark 6.7. The fact that every square in diagram (34) is c.i.p. means that

$$(36) \quad \mathcal{V}_0^{m-1} = \mathcal{V}_0^{m-2} \cap \mathcal{W}_0^m \quad \text{for any } 4 \leq m \leq n-1.$$

In particular Figure 2 represents geometrically the stratification of $\mathcal{V}_0^1 \cup \mathcal{W}_2^3 \subset T^1$ in the case of a D_6 singularity, by setting

$$F = \mathcal{V}_0^1 \cup \mathcal{W}_2^3, \quad E = \mathcal{V}_0^2, \quad D = \mathcal{W}_2^4, \quad C = \mathcal{V}_0^3, \quad B = \mathcal{W}_2^5, \quad A = \mathcal{V}_0^4 = \{0\}.$$

Proof. By the Morse Splitting Lemma 6.2, our problem can be reduced to the case $N = 1$ with $f(y, z) = y^2 z + z^{n-1}$ for $n \geq 4$. Let us start by typing

```
> for n from 4 to 10 do FD[n]:=y^2*z+z^(n-1) end do
      FD4 := y^2*z + z^3
      FD5 := y^2*z + z^4
      FD6 := y^2*z + z^5
      FD7 := y^2*z + z^6
      FD8 := y^2*z + z^7
      FD9 := y^2*z + z^8
      FD10 := y^2*z + z^9
> for n from 4 to 10 do TB[n]:=TyurinaBasis(FD[n]) end do
```

$$\begin{aligned}
TB_4 &:= [z^2, z, y, 1] \\
TB_5 &:= [z^3, z^2, z, y, 1] \\
TB_6 &:= [z^4, z^3, z^2, z, y, 1] \\
TB_7 &:= [z^5, z^4, z^3, z^2, z, y, 1] \\
TB_8 &:= [z^6, z^5, z^4, z^3, z^2, z, y, 1] \\
TB_9 &:= [z^7, z^6, z^5, z^4, z^3, z^2, z, y, 1] \\
TB_{10} &:= [z^8, z^7, z^6, z^5, z^4, z^3, z^2, z, y, 1]
\end{aligned}$$

In general, for $n \geq 4$, the Kuranishi space T^1 will be then given by

$$T^1 = \langle 1, y, z, \dots, z^{n-2} \rangle_{\mathbb{C}}.$$

Given $\Lambda = (\lambda_0, \lambda, \lambda_1, \dots, \lambda_{n-2}) \in T^1$, the associated small deformation of U_0 is

$$U_{\Lambda} = \{f_{\Lambda}(y, z) := f(y, z) + \lambda y + \sum_{i=0}^{n-2} \lambda_i z^i = 0\}.$$

A solution of the jacobian system of partials is then given by a solution $p_{\Lambda} = (y_{\Lambda}, z_{\Lambda})$ of the following polynomial system in $\mathbb{C}[\lambda][y, z]$

$$(37) \quad \begin{cases} 2yz + \lambda = 0 \\ (n-1)z^{n-2} + y^2 + \sum_{i=1}^{n-2} i\lambda_i z^{i-1} = 0 \end{cases}$$

giving precisely n critical points for f_{Λ} . Then p_{Λ} , one of them, is actually a singular point of U_{Λ} if and only if

$$(38) \quad p_{\Lambda} \in U_{\Lambda} \iff \Lambda \in \mathcal{L} := \{y_{\Lambda}^2 z_{\Lambda} + z_{\Lambda}^{n-1} + \lambda y_{\Lambda} + \sum_{i=0}^{n-2} \lambda_i z_{\Lambda}^i = 0\} \subset T^1$$

where, as above, $\mathcal{L}$ is one of the n hypersurfaces of T^1 parameterizing small deformations of $0 \in U_0$ to nodes. After translating $y \mapsto y + y_{\Lambda}, z \mapsto z + z_{\Lambda}$, we get

$$\begin{aligned}
f_{\Lambda}(y + y_{\Lambda}, z + z_{\Lambda}) = f &+ \left(y_{\Lambda}^2 z_{\Lambda} + z_{\Lambda}^{n-1} + \lambda y_{\Lambda} + \sum_{i=0}^{n-2} \lambda_i z_{\Lambda}^i \right) \\
(39) \quad &+ (2y_{\Lambda} z_{\Lambda} + \lambda) y \\
&+ \left((n-1)z_{\Lambda}^{n-2} + y_{\Lambda}^2 + \sum_{i=1}^{n-2} i\lambda_i z_{\Lambda}^{i-1} \right) z \\
&+ 2y_{\Lambda} yz + z_{\Lambda} y^2 \\
&+ \sum_{k=2}^{n-2} \left(\binom{n-1}{k} z_{\Lambda}^{n-1-k} + \sum_{i=k}^{n-2} \binom{i}{k} \lambda_i z_{\Lambda}^{i-k} \right) z^k
\end{aligned}$$

Then by (38) and (37) the origin (i.e. $p_{\Lambda} \in U_{\Lambda}$) is at least a node. We want at first recover, by means of a suitable algebraic stratification of the Kuranishi space T^1 , the upper row of the adjacency diagram (22). At this purpose consider the case $n = 6$:

$$> \quad T := \text{nops}(TB[6]);$$

```

> F[Lambda] := FD[6]+lambda[0]+lambda*y+
> sum(lambda[i-1]*TB[6][T-i], i = 2 .. T-1);
FΛ := y2z + z5 + λ0 + λy + λ1z + λ2z2 + λ3z3 + λ4z4

```

Compare $F\Lambda$ with f_Λ above.

```

> solve(diff(F[Lambda], y), diff(F[Lambda], z), [y, z]);
[[y = -1/2 * (lambda / (RootOf (lambda^2 + 20_Z^6 + 4 lambda_1_Z^2 + 8 lambda_2_Z^3 + 12 lambda_3_Z^4 + 16 lambda_4_Z^5))),
z = RootOf (lambda^2 + 20_Z^6 + 4 lambda_1_Z^2 + 8 lambda_2_Z^3 + 12 lambda_3_Z^4 + 16 lambda_4_Z^5)]]

```

Compare these solutions with (y_Λ, z_Λ) in (37). Set $K := y_\Lambda$ and $L := z_\Lambda$ then

```

> z := Z+L : y := Y+K :
> F[Lambda] := collect(F[Lambda], [Y, Z], 'distributed');
FΛ := LY2 + ZY2 + 2KYZ + (λ + 2KL)Y + Z5 + (5L + λ4)Z4 +
(10L2 + λ3 + 4λ4L)Z3 + (6λ4L2 + λ2 + 3λ3L + 10L3)Z2 +
(4λ4L3 + 5L4 + K2 + λ1 + 3λ3L2 + 2λ2L)Z +
K2L + λ0 + λK + λ4L4 + L5 + λ2L2 + λ1L + λ3L3

```

Compare $F\Lambda$ in the latter output with (39). As already done for A_n singularities in (33), let us call v_k the coefficients of singular deformations $F\Lambda$ of FD_6 . Precisely, let us type:

```

> SD := [Y*Z, Y^2, seq(Z^i, i = 2 .. 4)];
SD := [YZ, Y2, Z2, Z3, Z4]
> s := nops(SD);
> FL := Y^2*Z+Z^5+sum(v[i]*SD[i+1], i = 0 .. s-1);
FL := Y2Z + Z5 + v0YZ + v1Y2 + v2Z2 + v3Z3 + v4Z4
> MGB := MilnorGroebnerBasis(FL, {Y, Z});
MGB := {2YZ + v0Z + 2v1Y, Y2 + 5Z4 + v0Y + 2v2Z + 3v3Z2 + 4v4Z3,
4v2YZ - 3v0v3Z2 - 4v0v4Z3 - 5v0Z4 - v0Y2 + (-v02 + 4v2v1)Y}
> for i to nops(MGB) do LeadingTerm(MGB[i], tdeg_min(Y,Z))end do;
v0, Z
2v2, Z
-v02 + 4v2v1, Y

```

The previous output gives the couples (*leading coefficient*, *leading monomial*) of the three generators in the standard basis MGB of the jacobian ideal J_{FL} w.r.t. the l.m.o. $tdeg_min(Y, Z)$ ³:

- the possible relations on the coefficients v_k making a Milnor number growing up have then to be chosen among the three leading coefficients.

On the contrary of the A_n case, since both v_0 and v_2 multiply the same leading monomial Z , they have to be annihilated together to get a Milnor number increasing, describing a codimension 2 algebraic subset of $\mathcal{L}$. Actually it turns out to be an

³It is defined as the *opposite* of the reverse lexicographic g.m.o. on variables Y, Z i.e. it is a graduated and lexicographic l.m.o. for which $Y < Z$.

algebraic subset of the codimension 1 algebraic subset of $\mathcal{L}$ obtained by annihilating the third leading coefficient

$$(40) \quad \mathcal{W}_2 := \{4v_1v_2 - v_0^2 = 0\}$$

Notice that this is the same codimension 1 algebraic subset of $\mathcal{L}$ obtained by imposing the vanishing of the Hessian determinant $\det(\text{Hess}(0)) = 0$, in fact

```
> with(VectorCalculus): with(LinearAlgebra):
> H:=unapply(Hessian(FL, [Y, Z]), [Y, Z]):
> print(Hess(0)=H(0, 0), det(Hess(0))=Determinant(H(0,0)));
```

$$\text{Hess}(0) = \begin{bmatrix} 2v_1 & v_0 \\ v_0 & 2v_2 \end{bmatrix}, \quad \det(\text{Hess}(0)) = -v_0^2 + 4v_2v_1$$

Moreover if at least one of the three coefficients v_0, v_1, v_2 does not vanish then $\text{crk}(0) \leq 1$ and we are considering deformations of $0 \in f^{-1}(0)$ to simple A_m singularities, by Theorem 6.3(2). Introducing relation (40) in the deformation FL means to impose that

$$v_0YZ + v_1Y^2 + v_2Z^2 = (w_1Y + w_2Z)^2$$

where $w_i^2 = v_i$. Then

```
> F2:=Y^2*Z+Z^5+(w[1]*Y+w[2]*Z)^2+sum(v[j]*SD[j+1], j=3..s-1);
```

$$F2 := Y^2Z + Z^5 + (w_1Y + w_2Z)^2 + v_3Z^3 + v_4Z^4$$

```
> print(mu=MilnorNumber(F2,Y,Z), tau=TYURINANumber(F2,Y,Z));
```

$$\mu = 2, \tau = 2$$

implying, by Theorem 6.3(2), that $\mathcal{W}_2 \subset \mathcal{L}$ parameterizes small deformations of $0 \in f^{-1}(0)$ to a simple A_2 singularity or equivalently that $0 \in f_{\Lambda}^{-1}(0)$ is a simple A_2 singular point for generic $\Lambda \in \mathcal{W}_2$. As above, consider now the following standard basis

```
> MGB2 := MilnorGroebnerBasis(F2, Y, Z);
```

$$\begin{aligned} MGB2 := \{ & 2YZ + 2(w_1Y + w_2Z)w_1, Y^2 + 5Z^4 + 2(w_1Y + w_2Z)w_2 + 3v_3Z^2 + \\ & 4v_4Z^3, 6w_1w_2v_3Z^2Y + (-6v_3w_1^2 - 4w_2^2)ZY^2 + (-6v_3w_1^4 - 6w_1^2w_2^2)Y^2 + \\ & -10w_1^2w_2^2Z^4 - 8w_1^2w_2^2v_4Z^3 \} \end{aligned}$$

whose leading terms are listed as follows

```
> for i to nops(MGB2) do LeadingTerm(MGB2[i], tdeg_min(Y,Z)) end do;
```

$$2w_1w_2, Z$$

$$2w_2^2, Z$$

$$-6v_3w_1^4 - 6w_1^2w_2^2, Y^2$$

Observe that the vanishing of w_2 , hence v_2 , does not change the Milnor number since one get Y as leading monomial from the first generator. Then, as before, the interesting relation is given by the third leading coefficient, precisely: $v_1(v_1v_3 + v_2)$. Set

$$\mathcal{V}_1 := \{v_1 = 0\} \quad , \quad \mathcal{W}_3 := \{v_1v_3 + v_2 = 0\}$$

and observe that:

```
> w[1] := 0;
```

```
> print(mu=MilnorNumber(F2,Y,Z), tau=TyurinaNumber(F2,Y,Z));
```

$$\mu = 3, \tau = 3$$

On the other hand, by setting $w_2 = iw_1w_3$ with $w_i^2 = v_i$, one gets ⁴

```
> F3 := Z*Y^2+Z^5+v[1]*(Y+I*w[3]*Z)^2+w[3]^2*Z^3+v[4]*Z^4;
      F3 := ZY^2 + Z^5 + v1(Y + iw3Z)^2 + w3^2Z^3 + v4Z^4
> print(mu=MilnorNumber(F3,Y,Z),tau=TyurinaNumber(F3,Y,Z));
      mu = 3, tau = 3
```

Once more Theorem 6.3(2) gives that

- $0 \in f_\Lambda^{-1}(0)$ is a simple A_3 singularity for a generic Λ in

$$(41) \quad \mathcal{W}_2 \cap (\mathcal{V}_1 \cup \mathcal{W}_3) = (\mathcal{V}_0 \cap \mathcal{V}_1) \cup (\mathcal{W}_2 \cap \mathcal{W}_3) \subset \mathcal{W}_2 \subset \mathcal{L}$$

where $\mathcal{V}_k := \{v_k = 0\}$. Let us go on by considering the first connected component in (41) and by applying the same argument as above, precisely

```
> v[1] := 0;
> MGB3:=MilnorGroebnerBasis(F2,{Y,Z});
MGB3 := {2Y^3 + 10Z^4Y + 8Yv4Z^3, Y^2 + 5Z^4 + 2w2^2Z + 3v3Z^2 + 4v4Z^3, 2YZ}
> for i to nops(MGB3) do LeadingTerm(MGB3[i],tdeg_min(Y,Z))end do;
      2w2^2, Z
      2, YZ
      2, Y^3
```

Therefore the only interesting relation is given by the first leading coefficient, precisely $v_2 = 0$:

```
> w[2] := 0;
> print(mu=MilnorNumber(F2,{Y,Z}),tau=TyurinaNumber(F2,{Y,Z}));
      mu = 4, tau = 4
```

Since now $\text{crk}(0) = 2$, Theorem 6.3(3) gives that:

- $0 \in f_\Lambda^{-1}(0)$ is a simple D_4 singularity for a generic point $\Lambda \in \mathcal{V}_0 \cap \mathcal{V}_1 \cap \mathcal{V}_2$.

Let us now consider the second connected component $\mathcal{W}_2 \cap \mathcal{W}_3$ in (41). As before

```
> MGB3b := MilnorGroebnerBasis(F3,Y,Z);
MGB3b := {2YZ + 2v1(Y + iw3Z), Y^2 + 5Z^4 + 2iv1(Y + iw3Z)w3 + 3w3^2Z^2 +
      4v4Z^3, 10iw3^3v1^2Z^4 + (-8iw3v1v4 - 6iw3^3)Y^2Z^2 + (8w3^2 + 8v1v4)Y^3Z +
      -8w3^2v1v4Z^3Y + (8v1^2v4 + 8w3^2v1)Y^3}
> for i to nops(MGB3b) do LeadingTerm(MGB3b[i],tdeg_min(Y,Z))end do;
      2iv1w3, Z
      -2w3^2v1, Z
      8v1^2v4 + 8w3^2v1, Y^3
```

The only interesting relation is given by the last leading coefficient, precisely $v_1(v_1v_4 + v_3)$. Set then

$$\mathcal{W}_4 := \{v_1v_4 + v_3 = 0\}$$

⁴The reader may check that setting $w_2 = -iw_1w_3$ leads to the same results.

and introduce the new relation in the algebraic set $\mathcal{W}_2 \cap \mathcal{W}_3$ by intersecting it with $\mathcal{V}_1$ and $\mathcal{W}_4$ to get the following codimension 1 algebraic subset

$$(42) \quad (\mathcal{V}_1 \cap \mathcal{W}_2 \cap \mathcal{W}_3) \cup (\mathcal{W}_2 \cap \mathcal{W}_3 \cap \mathcal{W}_4) .$$

First of all observe that the first connected component in (42) reduces to the previous case since $\mathcal{V}_1 \cap \mathcal{W}_2 \cap \mathcal{W}_3 = \mathcal{V}_0 \cap \mathcal{V}_1 \cap \mathcal{V}_2$. Let us then consider the second connected component in (42) by introducing the relation $w_3 = iw_1w_4$, with $w_i^2 = v_i$, in $F3$. Then

```
> F4:=Z*Y^2+Z^5+(w[1]*Y-w[1]^2*w[4]*Z)^2-w[1]^2*w[4]^2*Z^3+
> w[4]^2*Z^4;
F4 := ZY^2 + Z^5 + (w1Y - w1^2w4Z)^2 - w1^2w4^2Z^3 + w4^2Z^4
> print(mu=MilnorNumber(F4,{Y,Z}),tau=TYURINANumber(F4,{Y,Z}));
mu = 4, tau = 4
```

Theorem 6.3(2) then gives that

- $0 \in f_{\Lambda}^{-1}(0)$ is a simple A_4 singularity for a generic $\Lambda \in \mathcal{W}_2 \cap \mathcal{W}_3 \cap \mathcal{W}_4$.

At the moment we have gotten the following chain of codimension 1 algebraic subsets

$$\begin{array}{ccccccc} \mathcal{L} & \longleftarrow & \mathcal{W}_2 & \longleftarrow & (\mathcal{V}_0 \cap \mathcal{V}_1) \cup (\mathcal{W}_2 \cap \mathcal{W}_3) & \longleftarrow & \mathcal{W}_2 \cap \mathcal{W}_3 \cap \mathcal{W}_4 \\ & & & & \uparrow & & \\ & & & & \mathcal{V}_0 \cap \mathcal{V}_1 \cap \mathcal{V}_2 & & \end{array}$$

representing the adjacency diagram

$$\begin{array}{ccccccc} A_1 & \longleftarrow & A_2 & \longleftarrow & A_3 & \longleftarrow & A_4 \\ & & & & \uparrow & & \\ & & & & D_4 & & \end{array}$$

Go on by considering the standard basis of $J_{f_{\Lambda}}$ when $\Lambda \in \mathcal{V}_0 \cap \mathcal{V}_1 \cap \mathcal{V}_2$:

```
> v[0]:=0 : v[1]:=0 : v[2]:=0 :
> MGB4 := MilnorGroebnerBasis(FL,{Y,Z})
MGB4 := {-2Y^3 - 10Z^4Y - 8Yv4Z^3, 2YZ, Y^2 + 5Z^4 + 3v3Z^2 + 4v4Z^3}
> for i to nops(MGB4) do LeadingTerm(MGB4[i],tdeg_min(Y,Z))end do;
2, YZ
-2, Y^3
3v3, Z^2
```

and then impose the further condition given by the latter leading coefficient, namely $v_3 = 0$:

```
> v[3] := 0 :
> print(mu=MilnorNumber(FL,{Y,Z}),tau=TYURINANumber(FL,{Y,Z}));
mu = 5, tau = 5
```

Theorem 6.3(3) then gives that

- $0 \in f_{\Lambda}^{-1}$ is a simple D_5 singularity for generic $\Lambda \in \bigcap_{k=0}^3 \mathcal{V}_k$.

On the other hand, for $\Lambda \in \mathcal{W}_2 \cap \mathcal{W}_3 \cap \mathcal{W}_4$ set:

```

> MGB4b := MilnorGroebnerBasis(F4, {Y, Z});
MGB4b := {2YZ + 2(w1Y - w1^2w4Z)w1, Y^2 + 5Z^4 - 2(w1Y - w1^2w4Z)w1^2w4 +
-3w1^2w4^2Z^2 + 4w4^2Z^3, (-8w1^2w4^4 - 10w1^4w4^2)Z^3Y^2 + (-10w4^2 - 10w1^2)ZY^4 +
-10w1^5w4^3Z^4Y + (-10w1^3w4 - 10w1w4^3)Y^3Z^2 + (-10w1^2w4^2 - 10w1^4)Y^4}
> for i to nops(MGB4b) do LeadingTerm(MGB4b[i], tdeg_min(Y, Z)) end do;
-2w1^3w4, Z
2w1^4w4^2, Z
-10w1^2w4^2 - 10w1^4, Y^4

```

As usual the only interesting relation is the last one, precisely $v_1(v_1 + v_4)$. Set $\mathcal{W}_5 := \{v_1 + v_4 = 0\}$ and intersect the algebraic subset defined by the latter relation with the second component in (42) to get

$$\begin{aligned}
 (\mathcal{V}_1 \cup \mathcal{W}_5) \cap (\mathcal{W}_2 \cap \mathcal{W}_3 \cap \mathcal{W}_4) &= (\mathcal{V}_1 \cap \mathcal{W}_2 \cap \mathcal{W}_3 \cap \mathcal{W}_4) \cup (\mathcal{W}_2 \cap \mathcal{W}_3 \cap \mathcal{W}_4 \cap \mathcal{W}_5) \\
 (43) \qquad \qquad \qquad &= (\mathcal{V}_0 \cap \mathcal{V}_1 \cap \mathcal{V}_2 \cap \mathcal{V}_3) \cup (\mathcal{W}_2 \cap \mathcal{W}_3 \cap \mathcal{W}_4 \cap \mathcal{W}_5)
 \end{aligned}$$

The first connected component in the right term gives the previous case just considered, then look at the second component by introducing the relation $w_4 = iw_1$ in F_4 :

```

> F5 := Z*Y^2 + Z^5 + v[1]*(Y - I*v[1]*Z)^2 + v[1]^2*Z^3 - v[1]*Z^4;
F5 := ZY^2 + Z^5 + v1(Y - iv1Z)^2 + v1^2Z^3 - v1Z^4
> print(mu=MilnorNumber(F5, {Z, Y}), tau=TyurinaNumber(F5, {Z, Y}));
mu = 5, tau = 5

```

and Theorem 6.3(2) allows to conclude that

- $0 \in f_\Lambda^{-1}$ is a simple A_5 singularity for generic $\Lambda \in \bigcap_{k=2}^5 \mathcal{W}_k$.

Since the algebraic subsets in (43) parameterizes 1-parameter deformations, the further and last step is clearly the trivial deformation given by $0 \in T^1$, completing the inclusions diagram as follows

$$(44) \quad \begin{array}{ccccccc}
 \mathcal{L} & \longleftarrow & \mathcal{W}_2 & \longleftarrow & \mathcal{V}_0^1 \cup \mathcal{W}_2^3 & \longleftarrow & \mathcal{W}_2^4 & \longleftarrow & \mathcal{W}_2^5 \\
 & & & & \uparrow & & \uparrow & & \uparrow \\
 & & & & \mathcal{V}_0^2 & \longleftarrow & \mathcal{V}_0^3 & \longleftarrow & \{0\}
 \end{array}$$

where $\mathcal{V}_a^b := \bigcap_{k=a}^b \mathcal{V}_k$ and $\mathcal{W}_a^b := \bigcap_{k=a}^b \mathcal{W}_k$. Diagram (44) gives a stratification, by nested algebraic subsets, of the Kuranishi space of a simple D_6 singular point, representing the following adjacency diagram

$$(45) \quad \begin{array}{ccccccc}
 A_1 & \longleftarrow & A_2 & \longleftarrow & A_3 & \longleftarrow & A_4 & \longleftarrow & A_5 \\
 & & & & \uparrow & & \uparrow & & \uparrow \\
 & & & & D_4 & \longleftarrow & D_5 & \longleftarrow & D_6
 \end{array}$$

and geometrically represented by Figure 2, as already observed in Remark 6.7.

The recursive structure is now sufficiently clear to pass at the general step for $n \geq 4$. Recalling f_Λ as written in (39) define

$$(46) \quad \begin{aligned} v_0 &:= 2y_\Lambda \\ v_1 &:= z_\Lambda \\ v_k &:= \binom{n-1}{k} z_\Lambda^{n-1-k} + \sum_{i=k}^{n-2} \binom{i}{k} \lambda_i z_\Lambda^{i-k}, \quad k = 2, \dots, n-2 \end{aligned}$$

and consider the associated codimension 1 sub-schemes of $\mathcal{L}$

$$(47) \quad \begin{aligned} \mathcal{V}_k &:= \{v_k = 0\}, \quad 0 \leq k \leq n-2 \\ \mathcal{W}_k &:= \begin{cases} \{4v_1v_2 - v_0^2 = 0\} & \text{for } k = 2 \text{ (vanishing of } \det(\text{Hess})) \\ \{v_1v_k + v_{k-1} = 0\} & \text{for } 3 \leq k \leq n-2 \\ \{v_1 + v_{n-2} = 0\} & \text{for } k = n-1 \end{cases} \end{aligned}$$

Then diagrams (44) and (45) generalizes to give diagrams (34) and (35) in the statement, respectively. Let us conclude by pointing out that in diagram (44), $\mathcal{V}_0^3 = \mathcal{V}_0^2 \cap \mathcal{W}_2^4$. This fact generalizes to diagram (34) giving the stated relations (36) between spaces of such a stratification. $\square$

6.4. Simple singularities of E_6 type.

Theorem 6.8. *Let T^1 be the Kuranishi space of a simple N -dimensional singular point $0 \in f^{-1}(0)$ with*

$$f(x_1, \dots, x_{N+1}) = \sum_{i=1}^{N-1} x_i^2 + x_N^3 + x_{N+1}^4$$

The subset of T^1 parameterizing small deformations of $0 \in f^{-1}(0)$ to a simple node is the union of 6 hypersurfaces. Moreover, calling $\mathcal{L}$ any of those hypersurfaces, there exists a stratification of nested algebraic subsets giving rise to the following sequence of inclusions and c.i.p. squares

$$(48) \quad \begin{array}{ccccccc} \mathcal{L} & \longleftarrow & \mathcal{W}_2 & \longleftarrow & \mathcal{W}_2^3 & \longleftarrow & \widetilde{\mathcal{W}}_2^4 \longleftarrow \widetilde{\mathcal{W}}_2^4 \cap \mathcal{V}_2 \\ & & & & \uparrow & & \uparrow \\ & & & & \mathcal{V}_0^2 & \longleftarrow & \mathcal{V} \cap \mathcal{V}_0^2 \\ & & & & \uparrow & & \uparrow \\ & & & & \{0\} & & \uparrow \end{array}$$

verifying the Arnol'd's adjacency diagram

$$(49) \quad \begin{array}{ccccccc} A_1 & \longleftarrow & A_2 & \longleftarrow & A_3 & \longleftarrow & A_4 \longleftarrow A_5 \\ & & & & \uparrow & & \uparrow \\ & & & & D_4 & \longleftarrow & D_5 \\ & & & & & & \uparrow \\ & & & & & & E_6 \end{array}$$

where

- $\mathcal{L}$ is the hypersurface of T^1 defined by equation (52), keeping in mind (51),
- $\mathcal{V}_0^m := \bigcap_{k=0}^m \mathcal{V}_k$ and $\mathcal{W}_2^m := \bigcap_{k=2}^m \mathcal{W}_k$ where $\mathcal{V}_k, \mathcal{W}_k$ are hypersurfaces of $\mathcal{L}$ defined by equations (53), (54) and (55),
- $\mathcal{V}$ is a hypersurfaces of $\mathcal{L}$ defined by equation (59),
- $\widetilde{\mathcal{W}}_2^4$ is a codimension 3 Zariski closed subset of $\mathcal{L}$ defined in (58).

Remark 6.9. Analogously to what observed in remark 6.7, the fact that every square in diagram (48) is c.i.p. means that

$$(50) \quad \widetilde{\mathcal{W}}_2^4 \cap \mathcal{V}_0^2 = \mathcal{V} \cap \mathcal{V}_0^2 \quad , \quad \left(\widetilde{\mathcal{W}}_2^4 \cap \mathcal{V}_2 \right) \cap (\mathcal{V} \cap \mathcal{V}_0^2) = \{0\}$$

In particular Figure 2 represents geometrically the stratification of $\mathcal{W}_2^3 \subset T^1$, by setting

$$F = \mathcal{W}_2^3 \quad , \quad E = \mathcal{V}_0^2 \quad , \quad D = \widetilde{\mathcal{W}}_2^4 \quad , \quad C = \mathcal{V} \cap \mathcal{V}_0^2 \quad , \quad B = \widetilde{\mathcal{W}}_2^4 \cap \mathcal{V}_2 \quad , \quad A = \{0\} \quad .$$

Proof. By the Morse Splitting Lemma 6.2, our problem can be reduced to the case $N = 1$ with $f(y, z) = y^3 + z^4$. To get an explicit basis of the Kuranishi space T^1 type

$$\begin{aligned} &> \text{F} := y^3 + z^4; \\ & \quad \quad \quad F := y^3 + z^4 \\ &> \text{TyB} := \text{TyurinaBasis(F)}; \\ & \quad \quad \quad \text{TyB} := [yz^2, z^2, yz, z, y, 1] \end{aligned}$$

Therefore

$$T^1 \cong \langle 1, y, z, yz, z^2, yz^2 \rangle_{\mathbb{C}} \quad .$$

Given $\Lambda = (\lambda_0, \lambda_1, \dots, \lambda_5) \in T^1$, the associated deformation of U_0 is

$$\begin{aligned} U_{\Lambda} &= \{f_{\Lambda}(y, z) = 0\} \quad \text{where} \\ f_{\Lambda}(y, z) &:= f(y, z) + \lambda_0 + \lambda_1 y + \lambda_2 z + \lambda_3 yz + \lambda_4 z^2 + \lambda_5 yz^2 \end{aligned}$$

A solution of the jacobian system of partials is then given by a solution $p_{\Lambda} = (y_{\Lambda}, z_{\Lambda})$ of the following polynomial system in $\mathbb{C}[\lambda][y, z]$

$$(51) \quad \begin{cases} 3y^2 + \lambda_1 + \lambda_3 z + \lambda_5 z^2 = 0 \\ 4z^3 + 2\lambda_4 z + \lambda_2 + y(\lambda_3 + 2\lambda_5 z) = 0 \end{cases}$$

giving precisely 6 critical points for f_{Λ} . Then p_{Λ} , one of them, is actually a singular point of U_{Λ} if and only if

$$(52) \quad p_{\Lambda} \in U_{\Lambda} \iff \Lambda \in \mathcal{L} \quad \text{where}$$

$$\mathcal{L} := \{y_{\Lambda}^3 + z_{\Lambda}^4 + \lambda_0 + \lambda_1 y_{\Lambda} + \lambda_2 z_{\Lambda} + \lambda_3 y_{\Lambda} z_{\Lambda} + \lambda_4 z_{\Lambda}^2 + \lambda_5 y_{\Lambda} z_{\Lambda}^2 = 0\} \subset T^1$$

which is one of the 6 hypersurfaces of T^1 parameterizing small deformations of $0 \in U_0$ to nodes. After translating $y \mapsto y + y_{\Lambda}, z \mapsto z + z_{\Lambda}$, we get

$$\begin{aligned} f_{\Lambda}(y + y_{\Lambda}, z + z_{\Lambda}) &= \\ &f(y, z) + (y_{\Lambda}^3 + z_{\Lambda}^4 + \lambda_0 + \lambda_1 y_{\Lambda} + \lambda_2 z_{\Lambda} + \lambda_3 y_{\Lambda} z_{\Lambda} + \lambda_4 z_{\Lambda}^2 + \lambda_5 y_{\Lambda} z_{\Lambda}^2) \\ &+ (3y_{\Lambda}^2 + \lambda_1 + \lambda_3 z_{\Lambda} + \lambda_5 z_{\Lambda}^2) y \\ &+ (4z_{\Lambda}^3 + 2\lambda_4 z_{\Lambda} + \lambda_2 + y_{\Lambda}(\lambda_3 + 2\lambda_5 z_{\Lambda})) z \\ &+ (\lambda_3 + 2\lambda_5 z_{\Lambda}) yz + 3y_{\Lambda} y^2 + (6z_{\Lambda}^2 + \lambda_4 + \lambda_5 y_{\Lambda}) z^2 \\ &+ 4z_{\Lambda} z^3 + \lambda_5 yz^2 \end{aligned}$$

as may be verified by setting $K := y_\Lambda, L := z_\Lambda$ and typing:

```
> F[Lambda] := F+sum(lambda[i]*TyB[T-i], i = 0 .. T-1)
      FL := y^3 + z^4 + lambda_0 + lambda_1 y + lambda_2 z + lambda_3 y z + lambda_4 z^2 + lambda_5 y z^2
> z:=Z+L : y:=Y+K :
> F[Lambda] := collect(F[Lambda], [Y, Z], 'distributed');
FL := Y^3 + 3 KY^2 + lambda_5 Y Z^2 + (lambda_3 + 2 lambda_5 L) Y Z + (lambda_3 L + lambda_1 + 3 K^2 + lambda_5 L^2) Y +
+ Z^4 + 4 L Z^3 + (lambda_4 + lambda_5 K + 6 L^2) Z^2 + (2 lambda_4 L + lambda_3 K + lambda_2 + 2 lambda_5 K L + 4 L^3) Z +
+ K^3 + lambda_0 + lambda_1 K + lambda_4 L^2 + lambda_2 L + L^4 + lambda_3 K L + lambda_5 K L^2
```

Define:

(53)

$$v_0 = \lambda_3 + 2\lambda_5 z_\Lambda, \quad v_1 = 3y_\Lambda, \quad v_2 = 6z_\Lambda^2 + \lambda_4 + \lambda_5 y_\Lambda, \quad v_3 = \lambda_5, \quad v_4 = 4z_\Lambda$$

and $\mathcal{V}_k := \{v_k = 0\}$. Then type:

```
> SD := [Y*Z, Y^2, Z^2, Y*Z^2, Z^3]:
> s := nops(SD):
> FL := Y^3+Z^4+sum(v[i]*SD[i+1], i = 0 .. s-1);
      FL := Y^3 + Z^4 + v_0 Y Z + v_1 Y^2 + v_2 Z^2 + v_3 Y Z^2 + v_4 Z^3
> MGB := MilnorGroebnerBasis(FL, {Y,Z});
MGB := { -6 v_2 Y^2 + (-2 v_2 v_3 + 3 v_0 v_4) Z^2 + 2 v_0 v_3 Y Z + (v_0^2 - 4 v_2 v_1) Y + 4 v_0 Z^3,
          3 Y^2 + v_0 Z + 2 v_1 Y + v_3 Z^2, 4 Z^3 + v_0 Y + 2 v_2 Z + 2 v_3 Y Z + 3 v_4 Z^2 }
> for i to nops(MGB) do LeadingTerm(MGB[i], tdeg_min(Y,Z)) end do;
      v_0^2 - 4 v_2 v_1, Y
      2 v_2, Z
      v_0, Z
```

The only interesting relation is given by the first leading coefficient giving the vanishing of $\det(\text{Hess}(0)) = 4v_1 v_2 - v_0^2$. Introduce it in FL by setting $v_k = w_k^2$ and considering

```
> F2:=Y^3+Z^4+(w[1]*Y+w[2]*Z)^2+sum(v[j]*SD[j+1], j = 3..s-1)
      F2 := Y^3 + Z^4 + (w_1 Y + w_2 Z)^2 + v_3 Y Z^2 + v_4 Z^3
> print(mu=MilnorNumber(F2,{Y,Z}), tau=TYURINANumber(F2,{Y,Z}));
      mu = 2, tau = 2
```

Then Theorem 6.3(2) ensures that

- $0 \in f_\Lambda^{-1}(0)$ is a simple A_2 singularity for Λ generic in

$$(54) \quad \mathcal{W}_2 := \{4v_1 v_2 - v_0^2 = 0\}.$$

Consider the standard basis:

```
> MGB2 := MilnorGroebnerBasis(F2, {Y,Z});
MGB2 := { 4 Z^3 + 2 (w_1 Y + w_2 Z) w_2 + 2 v_3 Y Z + 3 v_4 Z^2, (9 w_1^2 v_4 - 9 w_1 w_2 v_3) Y^3
          + (-6 w_2^3 w_1 + 6 w_1^4 v_4 - 6 w_1^3 w_2 v_3) Y^2 + (-3 w_1 w_2 v_3^2 + 3 w_1^2 v_4 v_3) Y Z^2
          + (-9 w_1 w_2 v_4 + 3 w_2^2 v_3) Y^2 Z + (8 w_1^2 w_2^2 - 3 v_4 v_3 w_1 w_2 + w_2^2 v_3^2) Z^3,
          3 Y^2 + 2 (w_1 Y + w_2 Z) w_1 + v_3 Z^2 }
```

```

> for i to nops(MGB2) do LeadingTerm(MGB2[i],tdeg_min(Y,Z))end do;
      2 w22, Z
      2 w1w2, Z
      -6 w23w1 + 6 w14v4 - 6 w13w2v3, Y2

```

Relations coming from the first and the second leading coefficients do not increase the Milnor number since:

```

> w[1] := 0:
> print(mu=MilnorNumber(F2,{Z,Y}),tau=TyurinaNumber(F2,{Z,Y}));
      μ = 2, τ = 2
> unassign('w[1]'): w[2] := 0 :
> print(mu=MilnorNumber(F2,{Z,Y}), tau=TYURINANumber(F2,{Z,Y}))
      μ = 2, τ = 2

```

Then consider the third leading coefficient, giving $v_4 = \frac{w_2}{w_1}v_3 + \left(\frac{w_2}{w_1}\right)^3$, since we can assume $w_1 \neq 0$. Then set

```

> w[2] := u*w[1] : v[4] := u*(v[3]+u^2) : F2 ;
      Y3 + Z4 + (w1Y + uw1Z)2 + v3YZ2 + u(v3 + u2)Z3
> F3 := Y^3+Z^4+v[1]*(Y+u*Z)^2+v[3]*Y*Z^2+u*(v[3]+u^2)*Z^3
      F3 := Y3 + Z4 + v1(Y + uZ)2 + v3YZ2 + u(v3 + u2)Z3
> print(mu=MilnorNumber(F3,{Z,Y}),tau=TYURINANumber(F3,{Z,Y}))
      μ = 3, τ = 3

```

and Theorem 6.3(2) gives that ⁵

• $0 \in f_{\Lambda}^{-1}(0)$ is a singularity of type A_3 for any generic $\Lambda \in \mathcal{W}_2 \cap \mathcal{W}_3$ where

$$(55) \quad \mathcal{W}_3 := \{v_1^3v_4^2 - v_2(v_1v_3 + v_2)^2 = 0\}.$$

The latter equation is obtained by observing that, after eliminating $w_1 \neq 0$, the third leading coefficient in $MGB2$ gives

$$w_1^3v_4 = \pm w_2(w_1^2v_3 + w_2^2)$$

(where the sign depends on the choice of the square roots $w_i^2 = v_i$) whose square is the equation of $\mathcal{W}_3$ in (55). Let us now consider the standard basis:

```

> MGB3 := MilnorGroebnerBasis(F3,{Y,Z});
MGB3 := { (16 v12 - 24 v1u2v3 - 4 v32v1 - 36 v1u4)Y3 + (18 u3v3 - 24 v1u + 6 uv32)ZY3 + (-3 u4v32 + 8 v1u2v3 - 2 u2v33)Z4 + (-36 u2v3 - 27 u4 - 6 v32 + 24 v1)Y4 + (8 v1v3 - 18 u2v32 + 24 v1u2 - 18 u4v3 - 2 v33)Z2Y2 + (-8 v1uv3 + 6 u3v32 + 2 uv33)Z3Y, 3 Y2 + 2 v1(Y + uZ) + v3Z2, 4 Z3 + 2 v1(Y + uZ)u + 2 v3YZ + 3 u(v3 + u2)Z2 }
> for i to nops(MGB3) do LeadingTerm(MGB3[i],tdeg_min(Y,Z))end do;
      2 v1u, Z
      2 v1u2, Z

```

⁵The reader may check that choosing $v_4 = -u(v_3 + u^2)$ leads to the same conclusion.

$$16 v_1^2 - 24 v_1 u^2 v_3 - 4 v_3^2 v_1 - 36 v_1 u^4, Y^3$$

The last leading coefficient can be rewritten as $v_1 [4v_1 - (v_3 + 3u^2)^2]$, giving all the new interesting relations, precisely $v_1 = 0$ and $v_1 = 1/4(v_3 + 3u^2)^2$. Since:

```
> v[1] := 0:
> print(mu=MilnorNumber(F3,{Z,Y}),tau=TyurinaNumber(F3,{Z,Y}))
      mu = 4, tau = 4
> unassign('v[1]') : v[1] := (1/4)*(3*u^2+v[3])^2 :
> print(mu=MilnorNumber(F3,{Z,Y}),tau=TyurinaNumber(F3,{Z,Y}))
      mu = 4, tau = 4
```

the Milnor number is then increasing for Λ in

$$(56) \quad (\mathcal{V}_1 \cap \mathcal{W}_2 \cap \mathcal{W}_3) \cup (\mathcal{W}_2 \cap \mathcal{W}_3 \cap \mathcal{W}_4) = \left(\bigcap_{k=0}^2 \mathcal{V}_k \right) \cup \left(\bigcap_{k=2}^4 \mathcal{W}_k \right)$$

where, recalling that $u^2 = v_2/v_1$,

$$(57) \quad \mathcal{W}_4 := \{4v_1^3 - (v_1 v_3 + 3v_2)^2 = 0\}.$$

As usual by now, call $\mathcal{V}_a^b := \bigcap_{k=a}^b \mathcal{V}_k$ and $\mathcal{W}_a^b := \bigcap_{k=a}^b \mathcal{W}_k$. Then (56) rewrites as $\mathcal{V}_0^2 \cup \mathcal{W}_2^4$. Notice that

$$\mathcal{V}_1^2 \subseteq \mathcal{W}_3^4 \implies \mathcal{V}_0^2 \subseteq \mathcal{W}_2^4.$$

Define the following codimension 1 Zariski closed subset ⁶ of $\mathcal{W}_2^3$

$$(58) \quad \widetilde{\mathcal{W}}_2^4 := \overline{\mathcal{W}_2^4 \setminus \mathcal{V}_0^2}.$$

Then Theorem 6.3(2) implies that

- $0 \in f_\Lambda^{-1}(0)$ is a singularity of type A_4 for any generic $\Lambda \in \widetilde{\mathcal{W}}_2^4$

and Theorem 6.3(3) ensures that

- $0 \in f_\Lambda^{-1}(0)$ is a singularity of type D_4 for any generic $\Lambda \in \mathcal{V}_0^2$.

Let us firstly consider the latter case:

```
> v[0]:=0 : v[1]:=0 : v[2]:=0 : FL ;
      Y^3 + Z^4 + v_3 Y Z^2 + v_4 Z^3
> print(mu=MilnorNumber(FL,{Y,Z}),tau=TyurinaNumber(FL,{Y,Z}))
      mu = 4, tau = 4
> MGB4 := MilnorGroebnerBasis(FL,{Y,Z})
MGB4 := {3 Y^2 + v_3 Z^2, 4 Z^3 + 2 v_3 Y Z + 3 v_4 Z^2, 9 v_4 Y^2 - 4 v_3 Z^3 - 2 v_3^2 Y Z,
      -36 v_4 Y v_3 Z^3 - 8 v_3^3 Z^4 + (12 v_3^3 + 81 v_4^2) Y^3}
> for i to nops(MGB4) do LeadingTerm(MGB4[i],tdeg_min(Y,Z))end do;
      v_3, Z^2
```

⁶ Although (58) is a topological definition, $\widetilde{\mathcal{W}}_2^4$ is clearly an algebraic subset of $\mathcal{L}$. Unfortunately it is not a complete intersection but the reader may obtain the (long) list of generators of its defining ideal by employing the MAPLE command `EliminationIdeal` in the `PolynomialIdeals` package. Not all the so listed generators are effectively necessary. The reader may compare them with those obtained by employing the MAPLE command `eliminate`, being careful with possible multiplicity and reducibility of generators.

$$\begin{aligned}
& 3v_4, Z^2 \\
& -2v_3^2, YZ \\
& 12v_3^3 + 81v_4^2, Y^3
\end{aligned}$$

The only interesting relation is given by the last leading coefficient since:

```

> v[4] := 0 :
> print(mu=MilnorNumber(FL,{Y,Z}),tau=TyurinaNumber(FL,{Y,Z}))
      mu = 4, tau = 4
> unassign('v[4]') : v[3]:=0 :
> print(mu=MilnorNumber(FL,{Y,Z}),tau=TyurinaNumber(FL,{Y,Z}))
      mu = 4, tau = 4
> unassign('v[3]') : v[3]:= -3*a^2 : v[4] := -2*a^3 : FL
      Y^3 + Z^4 - 3a^2YZ^2 - 2a^3Z^3
mu = MilnorNumber(FL,{Y,Z}), tau = TyurinaNumber(FL,{Y,Z})
      mu = 5, tau = 5

```

Therefore if

$$(59) \quad \mathcal{V} := \{4v_3^3 + 27v_4^2 = 0\}$$

then Theorem 6.3(3) allows to conclude that

- $0 \in f_{\Lambda}^{-1}(0)$ is a singularity of type D_5 for Λ generic in $\mathcal{V} \cap \mathcal{V}_0^2$.

Let us now consider $\Lambda \in \widetilde{\mathcal{W}}_2^4$. Then

```

> F3;
      Y^3 + Z^4 + 1/4 (3u^2 + v3)^2 (Y + uZ)^2 + v3YZ^2 + u(v3 + u^2)Z^3
> MGB4b:=MilnorGroebnerBasis(F3,{Y,Z}):
> for i to nops(MGB4b) do
>   LeadingTerm(MGB4b[i],tdeg_min(Y,Z))end do;
      1/2 (3u^2 + v3)^2 u, Z
      1/2 (3u^2 + v3)^2 u^2, Z
      12709329141645/256 u^2v3^2 + 38127987424935/256 u^6 + 38127987424935/256 u^4v3 +
      1412147682405/256 v3^3, Y^4
> CC:= [seq(LeadingCoefficient(MGB4b[i],tdeg_min(Y,Z)),
>   i=1..nops(MGB4b))]:
> factor(CC[3]);
      1412147682405/256 (3u^2 + v3)^3

```

The latter factorization clarifies that the third leading coefficient does not give any further relations w.r.t. those given by the previous leading coefficients, namely $v_2(3v_2 + v_1v_3) = 0$. Define

$$(60) \quad \mathcal{W}_5 := \{v_1v_3 + 3v_2 = 0\}$$

and observe that:

```
> u := 0 : F3
      Y^3 + Z^4 + 1/4 v_3^2 Y^2 + v_3 Y Z^2
> print(mu=MilnorNumber(F3,{Y,Z}),tau=TyurinaNumber(F3,{Y,Z}))
      mu = 5, tau = 5
```

implying, by Theorem 6.3(2), that

- $0 \in f_\Lambda^{-1}(0)$ is a singularity of type A_5 for a generic Λ in

$$\widetilde{\mathcal{W}}_2^4 \cap \mathcal{V}_2 = \mathcal{W} \cap \bigcap_{k=0}^2 \mathcal{V}_{2k}$$

where $\mathcal{W} := \{v_3^2 - 4v_1 = 0\}$.

On the other hand, notice that $\widetilde{\mathcal{W}}_2^4 \cap \mathcal{W}_5 = \mathcal{V} \cap \mathcal{V}_0^2$ giving the D_5 case previously analyzed, as may also be directly checked as follows

```
> unassign('u') : v[3] := -3*u^2 : F3
      Y^3 + Z^4 - 3 u^2 Y Z^2 - 2 u^3 Z^3
```

Since $(\mathcal{V} \cap \mathcal{V}_0^2) \cup (\mathcal{W}_2^4 \cap \mathcal{V}_2)$ parameterizes 1-parameter deformations, the further and last step is clearly the trivial deformation given by $0 \in T^1$, completing the inclusions diagram (48), giving a stratification, by nested algebraic subsets, of the Kuranishi space of a simple E_6 singular point and representing the adjacency diagram (49). At last, to prove (50), let us observe that

$$\widetilde{\mathcal{W}}_2^4 \cap \mathcal{V}_0^2 = \mathcal{V} \cap \mathcal{V}_0^2$$

since the common solutions of equations

$v_0 = v_1 = v_2 = 0$, $4v_1v_2 = v_0^2$, $v_1u^2 = v_2$, $v_4 = u(v_3 + u^2)$, $4v_1 = (v_3 + 3u^2)^2$ are given by $(v_0, v_1, v_2, v_3, v_4) = (0, 0, 0, -3u^2, -2u^3)$. Moreover

$$\left(\widetilde{\mathcal{W}}_2^4 \cap \mathcal{V}_2\right) \cap (\mathcal{V} \cap \mathcal{V}_0^2) = \left(\mathcal{W} \cap \bigcap_{k=0}^2 \mathcal{V}_{2k}\right) \cap \mathcal{V} \cap \mathcal{V}_2 = \{0\}$$

since 0 is the only common solution of equations

$$v_0 = v_2 = v_4 = 0, \quad 4v_1 = v_3^2, \quad 4v_3^3 + 27v_4^2 = 0.$$

□

6.5. Simple singularities of E_7 type.

Theorem 6.10. *Let T^1 be the Kuranishi space of a simple N -dimensional singular point $0 \in f^{-1}(0)$ with*

$$f(x_1, \dots, x_{N+1}) = \sum_{i=1}^{N-1} x_i^2 + x_N^3 + x_N x_{N+1}^3$$

The subset of T^1 parameterizing small deformations of $0 \in f^{-1}(0)$ to a simple node is the union of n hypersurfaces. Moreover, calling $\mathcal{L}$ any of those hypersurfaces,

there exists a stratification of nested algebraic subsets giving rise to the following sequence of inclusions, c.i.p. squares and a hinged union of c.i.p. squares

$$(61) \quad \begin{array}{ccccccccc} \mathcal{L} & \longleftarrow & \mathcal{W}_2 & \longleftarrow & \mathcal{W}_2^3 & \longleftarrow & \widetilde{\mathcal{W}}_2^4 & \longleftarrow & \widetilde{\mathcal{W}}_2^5 \cup \widetilde{\mathcal{W}}_2'^5 & \longleftarrow & \widetilde{\mathcal{W}}_2'^6 \\ & & \uparrow & & \uparrow & & \uparrow & & \uparrow & & \uparrow \\ & & \mathcal{V}_0^2 & \longleftarrow & \mathcal{V} \cap \mathcal{V}_0^2 & \longleftarrow & \mathcal{V}' \cap \mathcal{V}_0^2 & & & & \\ & & & & \uparrow & & \uparrow & & \uparrow & & \uparrow \\ & & & & \mathcal{V} \cap \mathcal{V}_0^3 & \longleftarrow & \{0\} & & & & \end{array}$$

verifying the Arnol'd's adjacency diagram

$$(62) \quad \begin{array}{ccccccccc} A_1 & \longleftarrow & A_2 & \longleftarrow & A_3 & \longleftarrow & A_4 & \longleftarrow & A_5 & \longleftarrow & A_6 \\ & & & & \uparrow & & \uparrow & & \uparrow & & \uparrow \\ & & & & D_4 & \longleftarrow & D_5 & \longleftarrow & D_6 & & \\ & & & & & & \uparrow & & \uparrow & & \\ & & & & & & E_6 & \longleftarrow & E_7 & & \end{array}$$

where

- $\mathcal{L}$ is the hypersurface of T^1 defined by equation (65), keeping in mind (64),
- $\mathcal{V}_0^m := \bigcap_{k=0}^m \mathcal{V}_k$ and $\mathcal{W}_2^m := \bigcap_{k=2}^m \mathcal{W}_k$ where $\mathcal{V}_k, \mathcal{W}_k$ are hypersurfaces of $\mathcal{L}$ defined by equations (66), (67), (70) and (73),
- $\mathcal{V}$ and $\mathcal{V}'$ are a hypersurfaces and a codimension 2 complete intersection in $\mathcal{L}$ defined by equations (68) and (69), respectively,
- $\widetilde{\mathcal{W}}_2^k$ and $\widetilde{\mathcal{W}}_2'^k$ are Zariski closed subsets of $\mathcal{L}$ defined by (72), (73), (74) and (75).

In particular complete intersection properties in diagram (61) are summarized by the following relations:

$$(63) \quad \begin{aligned} \widetilde{\mathcal{W}}_2^4 \cap \mathcal{V}_0^2 &= \mathcal{V} \cap \mathcal{V}_0^2, \quad \widetilde{\mathcal{W}}_2^5 \cap (\mathcal{V} \cap \mathcal{V}_0^2) = \mathcal{V}' \cap \mathcal{V}_0^2, \quad \widetilde{\mathcal{W}}_2'^5 \cap (\mathcal{V} \cap \mathcal{V}_0^2) = \mathcal{V} \cap \mathcal{V}_0^3 \\ (\mathcal{V}' \cap \mathcal{V}_0^2) \cap (\mathcal{V} \cap \mathcal{V}_0^3) &= \{0\} = \widetilde{\mathcal{W}}_2'^6 \cap (\mathcal{V}' \cap \mathcal{V}_0^2). \end{aligned}$$

Proof. By the Morse Splitting Lemma 6.2, our problem can be reduced to the case $N = 1$ with $f(y, z) = y^3 + yz^3$. To get an explicit basis of the Kuranishi space T^1 type

$$\begin{aligned} &> \text{F} := \mathbf{y}^{\wedge 3} + \mathbf{y} * \mathbf{z}^{\wedge 3}; \\ &> \text{TyB} := \text{TyurinaBasis}(\text{F}); \\ &\quad \text{TyB} := [z^4, z^3, z^2, yz, z, y, 1] \end{aligned}$$

Therefore $T^1 \cong \langle 1, y, z, yz, z^2, z^3, z^4 \rangle_{\mathbb{C}}$ and given $\Lambda = (\lambda_0, \lambda_1, \dots, \lambda_6) \in T^1$, the associated deformation of U_0 is

$$\begin{aligned} U_{\Lambda} &= \{f_{\Lambda}(y, z) = 0\} \quad \text{where} \\ f_{\Lambda}(y, z) &:= f(y, z) + \lambda_0 + \lambda_1 y + \lambda_2 z + \lambda_3 yz + \lambda_4 z^2 + \lambda_5 z^3 + \lambda_6 z^4. \end{aligned}$$

A solution of the jacobian system of partial derivatives is then given by a solution $p_\Lambda = (y_\Lambda, z_\Lambda)$ of the following polynomial system in $\mathbb{C}[\lambda][y, z]$

$$(64) \quad \begin{cases} 3y^2 + z^3 + \lambda_1 + \lambda_3 z \equiv 0 \\ 3yz^2 + \lambda_2 + \lambda_3 y + 2\lambda_4 z + 3\lambda_5 z^2 + 4\lambda_6 z^3 = 0 \end{cases}$$

giving precisely 7 critical points for f_Λ . Then p_Λ , one of them, is actually a singular point of U_Λ if and only if

$$(65) \quad p_\Lambda \in U_\Lambda \iff \Lambda \in \mathcal{L} \subset T^1 \quad \text{where} \\ \mathcal{L} := \{y_\Lambda^3 + y_\Lambda z_\Lambda^3 + \lambda_0 + \lambda_1 y_\Lambda + \lambda_2 z_\Lambda + \lambda_3 y_\Lambda z_\Lambda + \lambda_4 z_\Lambda^2 + \lambda_5 z_\Lambda^3 + \lambda_6 z_\Lambda^4 = 0\} \quad .$$

After translating $y \mapsto y + y_\Lambda, z \mapsto z + z_\Lambda$, we get

$$\begin{aligned} f_\Lambda(y + y_\Lambda, z + z_\Lambda) = \\ f(y, z) + (y_\Lambda^3 + y_\Lambda z_\Lambda^3 + \lambda_0 + \lambda_1 y_\Lambda + \lambda_2 z_\Lambda + \lambda_3 y_\Lambda z_\Lambda + \lambda_4 z_\Lambda^2 + \lambda_5 z_\Lambda^3 + \lambda_6 z_\Lambda^4) \\ + (3y_\Lambda^2 + z_\Lambda^3 + \lambda_1 + \lambda_3 z_\Lambda) y \\ + (3y_\Lambda z_\Lambda^2 + \lambda_2 + \lambda_3 y_\Lambda + 2\lambda_4 z_\Lambda + 3\lambda_5 z_\Lambda^2 + 4\lambda_6 z_\Lambda^3) z \\ + (3z_\Lambda^2 + \lambda_3) yz + 3y_\Lambda y^2 + (3y_\Lambda z_\Lambda + \lambda_4 + 3\lambda_5 z_\Lambda + 6\lambda_6 z_\Lambda^2) z^2 \\ + 3z_\Lambda yz^2 + (\lambda_5 + 4\lambda_6 z_\Lambda + y_\Lambda) z^3 + \lambda_6 z^4 \end{aligned}$$

as can be checked by setting $K := y_\Lambda, L := z_\Lambda$ and typing:

```
> T := nops(TyB):
> F[Lambda] := F+sum(lambda[i]*TyB[T-i], i= 0 .. T-1)
    F\Lambda := y^3 + yz^3 + \lambda_0 + \lambda_1 y + \lambda_2 z + \lambda_3 yz + \lambda_4 z^2 + \lambda_5 z^3 + \lambda_6 z^4
> z := Z+L : y := Y+K :
> F[Lambda] := collect(F[Lambda], [Y, Z],
> 'distributed');
    F\Lambda := Y^3 + 3KY^2 + YZ^3 + 3LYZ^2 + (\lambda_3 + 3L^2)YZ +
    (\lambda_3 L + 3K^2 + \lambda_1 + L^3)Y + \lambda_6 Z^4 + (\lambda_5 + K + 4\lambda_6 L)Z^3 +
    (\lambda_4 + 3KL + 3\lambda_5 L + 6\lambda_6 L^2)Z^2 +
    (3KL^2 + \lambda_2 + 4\lambda_6 L^3 + 2\lambda_4 L + \lambda_3 K + 3\lambda_5 L^2)Z +
    K^3 + \lambda_6 L^4 + \lambda_0 + \lambda_4 L^2 + KL^3 + \lambda_3 KL + \lambda_1 K + \lambda_2 L + \lambda_5 L^3
```

Define:

$$(66) \quad \begin{aligned} v_0 &= 3z_\Lambda^2 + \lambda_3, & v_1 &= 3y_\Lambda, & v_2 &= 3y_\Lambda z_\Lambda + \lambda_4 + 3\lambda_5 z_\Lambda + 6\lambda_6 z_\Lambda^2, \\ v_3 &= 3z_\Lambda, & v_4 &= \lambda_5 + 4\lambda_6 z_\Lambda + y_\Lambda, & v_5 &= \lambda_6, \end{aligned}$$

and $\mathcal{V}_k := \{v_k = 0\}$. Then the proof goes on exactly as in the E_6 case until D_4 singularities: precisely setting

$$(67) \quad \begin{aligned} \mathcal{W}_2 &= \{4v_1 v_2 - v_0^2 = 0\} \\ \mathcal{W}_3 &= \{v_1^3 v_4^2 - v_2(v_1 v_3 + v_2)^2 = 0\} \end{aligned}$$

we have the inclusions's chain

$$T^1 \supset \mathcal{L} \supset \mathcal{W}_2 \supset \mathcal{W}_2^3 \supset \mathcal{V}_1 \cap \mathcal{W}_2^3 = \mathcal{V}_0^2$$

of subsets parameterizing small deformations whose generic fibre is either smooth or admits a singularity of type A_1 , A_2 , A_3 or D_4 , respectively: this fact can be checked as follows ⁷

```

> SD := [Y*Z, Y^2, Z^2, Y*Z^2, Z^3, Z^4]:
> s := nops(SD):

FL := Y^3 + YZ^3 + v_0YZ + v_1Y^2 + v_2Z^2 + v_3YZ^2 + v_4Z^3 + v_5Z^4

> MGB:=MilnorGroebnerBasis(FL,{Y,Z},plex_min(Z,Y))
MGB := {3Y^2 + Z^3 + v_0Z + 2v_1Y + v_3Z^2, 6v_3Y^2Z + (4v_1v_2 - v_0^2)Z +
(8v_1v_5 - v_0)Z^3 + (-v_0v_3 + 6v_1v_4)Z^2 + 9Y^2Z^2 + 12Yv_5Z^3 + (9v_4 + 6v_1)YZ^2
+ (6v_2 + 4v_1v_3)YZ, 3YZ^2 + v_0Y + 2v_2Z + 2v_3YZ + 3v_4Z^2 + 4v_5Z^3}
> for i to nops(MGB) do LeadingTerm(MGB[i],plex_min(Z,Y))end do;
      v_0, Y
      2v_1, Y
      4v_1v_2 - v_0^2, Z
> v[0] := 0 :
> print(mu=MilnorNumber(FL,{Y,Z},plex_min(Z,Y)));
      mu = 1
> unassign('v[0]') : v[2] := 0 :
> print(mu=MilnorNumber(FL,{Y,Z},plex_min(Z,Y)));
      mu = 1
> unassign('v[2]') :
> F2:=Y^3+Y*Z^3+(w[1]*Y+w[2]*Z)^2+sum(v[j]*SD[j+1],j=3..s-1)
      F2 := Y^3 + YZ^3 + (w_1Y + w_2Z)^2 + v_3YZ^2 + v_4Z^3 + v_5Z^4
> print(mu=MilnorNumber(F2,{Y,Z},plex_min(Z,Y)))
      mu = 2
> MGB2:=MilnorGroebnerBasis(F2,{Y,Z},plex_min(Z,Y));
MGB2 := {(2w_1w_2v_3 + 12w_2^2v_5)Z^4 +
(2w_1v_3^2w_2 + 2w_2^2w_1^2 + 9w_2^2v_4 - 8w_2w_1^3v_5)Z^3 +
(6w_2^2w_1^2v_3 + 6w_2^4 - 6w_2w_1^3v_4)Z^2 - 9w_2w_1Y^2Z^2 + (9w_2^2 - 12w_2w_1v_5)YZ^3
+ (-9w_2w_1v_4 - 6w_2w_1^3 + 6w_2^2v_3)YZ^2, 3YZ^2 + 2(w_1Y + w_2Z)w_2 + 2v_3YZ
+ 3v_4Z^2 + 4v_5Z^3, 3Y^2 + Z^3 + 2(w_1Y + w_2Z)w_1 + v_3Z^2}
> for i to nops(MGB2) do LeadingTerm(MGB2[i],plex_min(Z,Y))end do;
      6w_2^2w_1^2v_3 + 6w_2^4 - 6w_2w_1^3v_4, Z^2
      2w_2w_1, Y
      2w_1^2, Y

```

⁷In the following computation we do not employ the default l.m.o. but `plex_min(Z,Y)`, which is the l.m.o. defined as the opposite of the pure lexicographic g.m.o. with $Y < Z$. In fact the latter l.m.o. turned out to be considerably more efficient than `tdeg_min(Y,Z)`.

```

> print(mu=MilnorNumber(F2,{Z,Y},plex_min(Z,Y)),
> tau=TYURINANumber(F2,{Z,Y},[plex_min(Z,Y),tdeg(Z,Y)]));
       $\mu = 2, \tau = 2$ 
> unassign('w[1]') : w[2]:=0 :
> print(mu=MilnorNumber(F2,{Y,Z},plex_min(Z,Y)),
> tau=TyurinaNumber(F2,{Y,Z},plex_min(Z,Y)))
       $\mu = 2, \tau = 2$ 
> unassign('w[2]') : w[2] := u*w[1] : v[4] := u*(v[3]+u^2) :
> F2 ;
       $Y^3 + YZ^3 + (w_1Y + uw_1Z)^2 + v_3YZ^2 + u(v_3 + u^2)Z^3 + v_5Z^4$ 
> F3:=Y^3+Y*Z^3+v[1]*(Y+u*Z)^2+v[3]*Y*Z^2+u*(v[3]+u^2)*Z^3
> +v[5]*Z^4
       $F3 := Y^3 + YZ^3 + v_1(Y + uZ)^2 + v_3YZ^2 + u(v_3 + u^2)Z^3 + v_5Z^4$ 
> print(mu=MilnorNumber(F3,{Y,Z},plex_min(Z,Y)),
> tau=TYURINANumber(F3,{Y,Z},[plex_min(Z,Y),tdeg(Z,Y)]))
       $\mu = 3, \tau = 3$ 
> MGB3:=MilnorGroebnerBasis(F3,{Y,Z},plex_min(Z,Y));
       $MGB3 := \{ (-27u^6 - 9u^2v_3^2 - 10v_1uv_3 - 24v_1u^2v_5 - 36u^4v_3)Z^4 +$ 
       $(-18v_1u^2 + 24v_1uv_5 - 6uv_3^2 - 18u^3v_3)YZ^3 +$ 
       $(-24v_1v_3u^3 + 16uv_1^2v_5 - 36u^5v_1 - 4v_1uv_3^2 - 16v_1^2u^2)Z^3 +$ 
       $(-9uv_3 - 27u^3)YZ^4 + (-36u^3v_5 - 6v_1u - 12uv_3v_5)Z^5,$ 
       $3YZ^2 + 2v_1(Y + uZ)u + 2v_3YZ + 3u(v_3 + u^2)Z^2 + 4v_5Z^3,$ 
       $3Y^2 + Z^3 + 2v_1(Y + uZ) + v_3Z^2 \}$ 
> for i to nops(MGB3) do LeadingTerm(MGB3[i],plex_min(Z,Y))end do;
       $-24v_1v_3u^3 + 16uv_1^2v_5 - 36u^5v_1 - 4v_1uv_3^2 - 16v_1^2u^2, Z^3$ 
       $2v_1u, Y$ 
       $2v_1, Y$ 
> u := 0 :
> print(mu=MilnorNumber(F3,{Z,Y},plex_min(Z,Y)),
> tau = TyurinaNumber(F3,{Z,Y},plex_min(Z,Y)))
       $\mu = 3, \tau = 3$ 
> unassign('u') : v[1] := 0 :
> print(mu=MilnorNumber(F3,{Y,Z},plex_min(Z,Y)),
> tau=TyurinaNumber(F3,{Y,Z},plex_min(Z,Y)))
       $\mu = 4, \tau = 4$ 

```

Also the singularities' specialization in fibers parameterized by $\mathcal{V}_0^2$ proceeds as in the case of E_6 singularities. Precisely:

```

> v[0] := 0 : v[1] := 0 : v[2] := 0 : FL
       $Y^3 + YZ^3 + v_3YZ^2 + v_4Z^3 + v_5Z^4$ 
> MGB4 := MilnorGroebnerBasis(FL,{Z,Y},plex_min(Z,Y))
       $MGB4 := \{ 3Y^2 + Z^3 + v_3Z^2, (10v_3^2 + 36v_4v_5)Z^4 + (-24v_3v_5 + 27v_4)YZ^3$ 
       $+ (4v_3^3 + 27v_4^2)Z^3 + 6v_3Z^5, 3YZ^2 + 2v_3YZ + 3v_4Z^2 + 4v_5Z^3 \}$ 

```

```

> for i to nops(MGB4) do LeadingTerm(MGB4[i],plex_min(Z,Y))end do;
      4 v33 + 27 v42, Z3
      3, Y2
      2 v3, YZ
> v[3] := 0 :
> print(mu=MilnorNumber(FL,{Z,Y},plex_min(Z,Y)),
> tau=TyurinaNumber(FL,{Z,Y},plex_min(Z,Y)))
      μ = 4, τ = 4
> unassign('v[3]') : v[3] := -3*a^2 : v[4] := -2*a^3 : FL
      Y3 + YZ3 - 3 a2YZ2 - 2 a3Z3 + v5Z4
> print(mu=MilnorNumber(FL,{Z,Y},plex_min(Z,Y)),
> tau=TyurinaNumber(FL,{Z,Y},plex_min(Z,Y)))
      μ = 5, τ = 5

```

Therefore if

$$(68) \quad \mathcal{V} := \{4v_3^3 + 27v_4^2 = 0\}$$

then Theorem 6.3(3) allows to conclude that

- $0 \in f_{\Lambda}^{-1}(0)$ is a singularity of type D_5 for Λ generic in $\mathcal{V} \cap \mathcal{V}_0^2$.

Moreover:

```

> MGB5 := MilnorGroebnerBasis(FL,{Z,Y},plex_min(Z,Y));
MGB5 := { (48 a4 - 48 a3v5) Z4 + (12 v5 - 9 a) Z4Y + (16 v52 - 12 av5 - 6 a2) Z5,
          3 YZ2 - 6 a2YZ - 6 a3Z2 + 4 v5Z3, 3 Y2 + Z3 - 3 a2Z2 }
> for i to nops(MGB5) do LeadingTerm(MGB5[i],plex_min(Z,Y))end do;
      48 a4 - 48 a3v5, Z4
      -6 a2, YZ
      3, Y2
> a := 0 : FL
      Y3 + YZ3 + v5Z4
> print(mu=MilnorNumber(FL,{Z,Y},plex_min(Z,Y)),
> tau=TyurinaNumber(FL,{Z,Y},plex_min(Z,Y)))
      μ = 6, τ = 6
> unassign('a') : v[5] := a : FL
      Y3 + YZ3 - 3 a2YZ2 - 2 a3Z3 + aZ4
> print(mu=MilnorNumber(FL,{Z,Y},plex_min(Z,Y)),
> tau=TyurinaNumber(FL,{Z,Y},plex_min(Z,Y)))
      μ = 6, τ = 6

```

Then Theorem 6.3(4) gives that

- $0 \in f_{\Lambda}^{-1}(0)$ is a singularity of type E_6 for Λ generic in $\mathcal{V} \cap \mathcal{V}_0^3 = \mathcal{V}_0^4$.

On the other hand, by defining the following codimension 2 subset of $\mathcal{L}$

$$(69) \quad \mathcal{V}' = \{v_3 + 3v_5^2 = v_4 + 2v_5^3 = 0\} ,$$

Theorem 6.3(3) gives that

- $0 \in f_{\Lambda}^{-1}(0)$ is a singularity of type D_6 for Λ generic in $\mathcal{V}' \cap \mathcal{V}_0^2$

A further specialization here gives then the trivial deformation since $(\mathcal{V} \cap \mathcal{V}_0^3) \cap (\mathcal{V}' \cap \mathcal{V}_0^2) = \mathcal{V}' \cap \mathcal{V}_0^3 = \{0\}$.

Let us come back to consider $\mathcal{W}_2^3$ and look at the first leading coefficient in *MGB3*, giving the relation

$$4v_1(v_5 - u) - (v_3 + 3u^2)^2 = 0.$$

Then set $v_5 := t^2 + u$ and $v_3 := \pm 2w_1t - 3u^2$ and type ⁸

```
> v[3] := 2*t*w[1] - 3*u^2 : v[5] := t^2 + u : F2
      Y^3 + YZ^3 + (w1Y + uw1Z)^2 +
      (2tw1 - 3u^2)YZ^2 + u(2tw1 - 2u^2)Z^3 + (t^2 + u)Z^4
> print(mu=MilnorNumber(F2,{Y,Z},plex_min(Z,Y)),
> tau = TYURINANumber(F2,{Y,Z},[plex_min(Z,Y),tdeg(Z,Y)]))
      mu = 4, tau = 4
```

Define

$$(70) \quad \mathcal{W}_4 := \{16v_1^5v_2 - [(v_1v_2 + 3v_2)^2 - 4v_1^3v_5]^2 = 0\}$$

whose equation is obtained by eliminating u and t from the following set of equations, parameterizing $\mathcal{W}_3^4 \setminus \mathcal{V}_1^2$,

$$(71) \quad v_2 = u^2v_1, \quad v_4 = u(v_3 + u^2), \quad v_5 = u + t^2, \quad v_1 = w_1^2, \quad v_3 = \pm 2w_1t - 3u^2.$$

Observe that $\mathcal{V}_1^2 \subseteq \mathcal{W}_3^4 \Rightarrow \mathcal{V}_0^2 \subseteq \mathcal{W}_2^4$ and define the following codimension 1 Zariski closed subset ⁹ of $\mathcal{W}_2^3$

$$(72) \quad \widetilde{\mathcal{W}}_2^4 := \overline{\mathcal{W}_2^4 \setminus \mathcal{V}_0^2}.$$

Then Theorem 6.3(2) implies that

- $0 \in f_{\Lambda}^{-1}(0)$ is a singularity of type A_4 for any generic $\Lambda \in \widetilde{\mathcal{W}}_2^4$.

Moreover:

```
> MGB4b := MilnorGroebnerBasis(F2,{Y,Z},plex_min(Z,Y));
MGB4b := {3Y^2 + Z^3 + 2(w1Y + uw1Z)w1 + (2tw1 - 3u^2)Z^2,
3YZ^2 + 2(w1Y + uw1Z)uw1 + 2(2tw1 - 3u^2)YZ + 3u(2tw1 - 2u^2)Z^2 +
4(t^2 + u)Z^3, (-9w1 - 54ut)YZ^5 + (-9uw1 - 12w1t^2 - 72ut^3 - 54u^2t)Z^6 +
(-20tw1^4 - 80w1^3ut^2 - 60u^2t^3w1^2)Z^4 +
(-30tw1^2 - 102w1ut^2 + 90u^3t - 72u^2t^3 + 15w1u^2)YZ^4 +
(-24w1^2t^3 + 90u^4t - 38w1^2ut + 72u^3t^3 + 15u^3w1 - 60w1u^2t^2 - 24w1ut^4 - 6w1^3)Z^5}
> for i to nops(MGB4b) do LeadingTerm(MGB4b[i],plex_min(Z,Y))end do;
      2w1^2, Y
      2uw1^2, Y
      -20tw1^4 - 80w1^3ut^2 - 60u^2t^3w1^2, Z^4
> u := 0 : F2
      Y^3 + YZ^3 + w1^2Y^2 + 2tw1YZ^2 + t^2Z^4
```

⁸The reader may check that choosing $v_3 := -2w_1t - 3u^2$ gives the same result.

⁹The same considerations explained by footnote 6 are still holding, here.

```

> print(mu=MilnorNumber(F2,{Z,Y},plex_min(Z,Y)),
> tau=TYURINANumber(F2,{Z,Y},[plex_min(Z,Y), tdeg(Z,Y)]))
       $\mu = 4, \tau = 4$ 
> unassign('u') : w[1] := 0 : F2
       $Y^3 + YZ^3 - 3u^2YZ^2 - 2u^3Z^3 + (t^2 + u)Z^4$ 
> print(mu=MilnorNumber(F2,{Z,Y},plex_min(Z,Y)),
> tau=TyurinaNumber(F2,{Z,Y},plex_min(Z,Y)))
       $\mu = 5, \tau = 5$ 
> unassign('w[1]') : t:=0 : F2
       $Y^3 + YZ^3 + (w_1Y + uw_1Z)^2 - 3u^2YZ^2 - 2u^3Z^3 + Z^4u$ 
> print(mu=MilnorNumber(F2,{Z,Y},plex_min(Z,Y)),
> tau=TYURINANumber(F2,{Z,Y},[plex_min(Z,Y), tdeg(Z,Y)]))
       $\mu = 5, \tau = 5$ 

```

Then $\widetilde{\mathcal{W}}_2^4 \cap \mathcal{V}_1 = \mathcal{V} \cap \mathcal{V}_0^2$ reduces to the already considered case of generic D_5 singularities. On the other hand setting $t = 0$ means defining

$$(73) \quad \mathcal{W}_5 := \{v_1v_5^2 - v_2 = 0\} \quad , \quad \widetilde{\mathcal{W}}_2^5 := \overline{\mathcal{W}_2^5 \setminus \mathcal{V}_0^2} \quad ,$$

and observing that, by Theorem 6.3(2),

- $0 \in f_\Lambda^{-1}(0)$ is a singularity of type A_5 for generic $\Lambda \in \widetilde{\mathcal{W}}_2^4 \cap \mathcal{W}_5 = \widetilde{\mathcal{W}}_2^5$.

Consider then

```

> MGB5b := MilnorGroebnerBasis(F2,{Z,Y},plex_min(Z,Y));
       $MGB5b := \{3Y^2 + Z^3 + 2(w_1Y + uw_1Z)w_1 - 3u^2Z^2,$ 
       $-12w_1^4Z^5 + 12w_1^2Z^6u - 18w_1^2Z^5Y,$ 
       $3YZ^2 + 2(w_1Y + uw_1Z)uw_1 - 6YZu^2 - 6u^3Z^2 + 4Z^3u\}$ 
> for i to nops(MGB5b) do LeadingTerm(MGB5b[i],plex_min(Z,Y))end do;
       $2w_1^2, Y$ 
       $-12w_1^4, Z^5$ 
       $2uw_1^2, Y$ 
> u := 0 : F2
       $Y^3 + YZ^3 + w_1^2Y^2$ 
> print(mu=MilnorNumber(F2,{Z,Y},plex_min(Z,Y)),
> tau=TyurinaNumber(F2,{Z,Y},plex_min(Z,Y)))
       $\mu = 5, \tau = 5$ 
> unassign('u') : w[1]:=0 : F2
       $Y^3 + YZ^3 - 3u^2YZ^2 - 2u^3Z^3 + Z^4u$ 
> print(mu=MilnorNumber(F2,{Z,Y},plex_min(Z,Y)),
> tau=TyurinaNumber(F2,{Z,Y},plex_min(Z,Y)))
       $\mu = 6, \tau = 6$ 

```

which means that $\mathcal{V}_1 \cap \widetilde{\mathcal{W}}_2^5 = \mathcal{V}' \cap \mathcal{V}_0^2$ reduces to the already considered case of generic D_6 singularities.

Let us then come back to $\widetilde{\mathcal{W}}_2^4$ and consider the last leading coefficient in the standard basis $MGB4b$ giving the further relations $w_1 + ut = 0$ and $w_1 + 3ut = 0$, not yet analyzed. Then

```

> F2
      Y3 + YZ3 + (w1Y + uw1Z)2 + (2tw1 - 3u2)YZ2
      + u(2tw1 - 2u2)Z3 + (t2 + u)Z4

> w[1] := -u*t : F2
      Y3 + YZ3 + (-tuY - u2tZ)2 + (-2t2u - 3u2)YZ2
      + u(-2t2u - 2u2)Z3 + (t2 + u)Z4

> print(mu=MilnorNumber(F2,{Z,Y},plex_min(Z,Y)),
> tau=TYURINANumber(F2,{Z,Y},[plex_min(Z,Y),tdeg(Z,Y)]))
      μ = 4, τ = 4

> w[1] := -3*u*t : F2
      Y3 + YZ3 + (-3tuY - 3u2tZ)2 + (-6t2u - 3u2)YZ2
      + u(-6t2u - 2u2)Z3 + (t2 + u)Z4

> print(mu=MilnorNumber(F2,{Z,Y},plex_min(Z,Y)),
> tau=TYURINANumber(F2,{Z,Y},[plex_min(Z,Y),tdeg(Z,Y)]))
      μ = 5, τ = 5

```

and, by eliminating u, t and w_1 from equations $w_1 = -3ut$ and (71), define

$$(74) \quad \mathcal{W}'_5 := \{v_1(v_1^2 - 9v_2v_5)^2 - 81v_2^3 = 0\} \quad , \quad \widetilde{\mathcal{W}}'_2{}^5 := \widetilde{\mathcal{W}}_2^4 \cap \mathcal{W}'_5 .$$

Hence Theorem 6.3(2) gives that

- $0 \in f_\Lambda^{-1}(0)$ is a singularity of type A_5 for generic $\Lambda \in \widetilde{\mathcal{W}}'_2{}^5$.

Consider the associated standard basis:

```

> MGB5c:=MilnorGroebnerBasis(F2,{Z,Y},plex_min(Z,Y))
MGB5c := {3YZ2 - 6(-3tuY - 3u2tZ)u2t + 2(-6t2u - 3u2)YZ
+ 3u(-6t2u - 2u2)Z2 + 4(t2 + u)Z3,
3Y2 + Z3 - 6(-3tuY - 3u2tZ)tu + (-6t2u - 3u2)Z2,
(48t4 - 24t2u)Z5Y + (192t6u2 - 144u3t4)Z5 + (-16t4u + 16u2t2 + 64t6)Z6 }

> for i to nops(MGB5c) do LeadingTerm(MGB5c[i],plex_min(Z,Y))end do;
      18u3t2, Y
      18u2t2, Y
      192t6u2 - 144u3t4, Z5

> u := 0 : F2
      Y3 + YZ3 + t2Z4

> print(mu=MilnorNumber(F2,{Z,Y},plex_min(Z,Y)),
> tau=TYURINANumber(F2,{Z,Y},[plex_min(Z,Y),tdeg(Z,Y)]))
      μ = 6, τ = 6

> unassign('u') : t := 0 : F2
      Y3 + YZ3 - 3u2YZ2 - 2u3Z3 + Z4u

```

This gives precisely the same situation of generic E_6 and D_6 singularities previously considered, meaning that

- $0 \in f_\Lambda^{-1}(0)$ is a singularity of type E_6 for generic $\Lambda \in \mathcal{V}_2 \cap \widetilde{\mathcal{W}}_2^5 = \mathcal{V}_0^4$,
- $0 \in f_\Lambda^{-1}(0)$ is a singularity of type D_6 for generic $\Lambda \in \mathcal{W}_5 \cap \widetilde{\mathcal{W}}_2^5 = \widetilde{\mathcal{W}}_2^5 \cap \mathcal{W}_5' = \mathcal{V}' \cap \mathcal{V}_0^2$.

Moreover the last leading coefficient in $MGB5c$ gives the further relation

```
> u := (4/3)*t^2 : F2
Y^3 + YZ^3 + (-4t^3Y - 16/3t^5Z)^2 - 40/3t^4YZ^2 - 416/27t^6Z^3 + 7/3t^2Z^4

> print(mu=MilnorNumber(F2,{Z,Y},plex_min(Z,Y)),
> tau=TYURINANumber(F2,{Z,Y},[plex_min(Z,Y),tdeg(Z,Y)]))
mu = 6, tau = 6
```

meaning that

- $0 \in f_\Lambda^{-1}(0)$ is a singularity of type A_6 for generic $\Lambda \in \widetilde{\mathcal{W}}_2^6$

where

$$(75) \quad \widetilde{\mathcal{W}}_2^6 := \widetilde{\mathcal{W}}_2^5 \cap \mathcal{W}_6 \quad \text{and} \quad \mathcal{W}_6 := \{16v_1^5 - 729v_2^3 = 0\}$$

The reader can then easily check that (63) holds. $\square$

6.6. Simple singularities of E_8 type.

Theorem 6.11. *Let T^1 be the Kuranishi space of a simple N -dimensional singular point $0 \in f^{-1}(0)$ with*

$$f(x_1, \dots, x_{N+1}) = \sum_{i=1}^{N-1} x_i^2 + x_N^3 + x_{N+1}^5$$

The subset of T^1 parameterizing small deformations of $0 \in f^{-1}(0)$ to a simple node is the union of n hypersurfaces. Moreover, calling $\mathcal{L}$ any of those hypersurfaces, there exists a stratification of nested algebraic subsets giving rise to the following sequence of inclusions, c.i.p. squares and reducible c.i.p. squares

$$(76) \quad \begin{array}{ccccccccccc} \mathcal{L} & \longleftarrow & \mathcal{W}_2 & \longleftarrow & \mathcal{W}_2^3 & \longleftarrow & \widetilde{\mathcal{W}}_2^4 & \longleftarrow & \widetilde{\mathcal{W}}_2^5 & \longleftarrow & \widetilde{\mathcal{W}}_2^6 & \longleftarrow & \widetilde{\mathcal{W}}_2^7 \\ & & & & \uparrow & & \uparrow & & \uparrow & & \uparrow & & \uparrow \\ & & & & \mathcal{V}_0^2 & \longleftarrow & \mathcal{V} \cap \mathcal{V}_0^2 & \longleftarrow & \mathcal{V}' \cap \mathcal{V}_0^2 & \longleftarrow & \mathcal{V}'' \cap \mathcal{V}_0^2 & & \\ & & & & & & \uparrow & & \uparrow & & \uparrow & & \uparrow \\ & & & & & & \mathcal{V}_0^4 & \longleftarrow & \mathcal{V}_0^4 \cap \mathcal{V}_6 & \longleftarrow & \{0\} & & \end{array}$$

verifying the Arnol'd's adjacency diagram

$$(77) \quad \begin{array}{cccccccc} A_1 & \longleftarrow & A_2 & \longleftarrow & A_3 & \longleftarrow & A_4 & \longleftarrow & A_5 & \longleftarrow & A_6 & \longleftarrow & A_7 \\ & & & & \uparrow & & \uparrow & & \uparrow & & \uparrow & & \uparrow \\ & & & & D_4 & \longleftarrow & D_5 & \longleftarrow & D_6 & \longleftarrow & D_7 & & \\ & & & & & & \uparrow & & \uparrow & & \uparrow & & \\ & & & & & & E_6 & \longleftarrow & E_7 & \longleftarrow & E_8 & & \end{array}$$

where

- $\mathcal{L}$ is the hypersurface of T^1 defined by equation (80), keeping in mind (79),
- $\mathcal{V}_0^m := \bigcap_{k=0}^m \mathcal{V}_k$ and $\mathcal{W}_2^m := \bigcap_{k=2}^m \mathcal{W}_k$ where $\mathcal{V}_k, \mathcal{W}_k$ are hypersurfaces of $\mathcal{L}$ defined by equations (82), (83), (86), (90), (92) and (93),
- $\mathcal{V}, \mathcal{V}'$ and $\mathcal{V}''$ are a hypersurface, a codimension 2 and a codimension 3 complete intersections in $\mathcal{L}$, respectively, defined by the latter equation in (83), by (84) and by (85), respectively,
- $\widetilde{\mathcal{W}}_2^k$ are Zariski closed subsets of $\mathcal{L}$ defined by (88), (89), (91) and (93).

In particular, complete intersection properties in diagram (76) are summarized by the following relations:

$$(78) \quad \begin{aligned} \widetilde{\mathcal{W}}_2^4 \cap \mathcal{V}_0^2 &= \mathcal{V} \cap \mathcal{V}_0^2, \\ \widetilde{\mathcal{W}}_2^5 \cap (\mathcal{V} \cap \mathcal{V}_0^2) &= (\mathcal{V}' \cap \mathcal{V}_0^2) \cup \mathcal{V}_0^4, \\ \widetilde{\mathcal{W}}_2^6 \cap (\mathcal{V}' \cap \mathcal{V}_0^2) &= (\mathcal{V}'' \cap \mathcal{V}_0^2) \cup (\mathcal{V}_0^4 \cap \mathcal{V}_6), \\ (\mathcal{V}'' \cap \mathcal{V}_0^2) \cap (\mathcal{V}_0^4 \cap \mathcal{V}_6) &= \widetilde{\mathcal{W}}_2^7 \cap (\mathcal{V}'' \cap \mathcal{V}_0^2) = \{0\}. \end{aligned}$$

Proof. By the Morse Splitting Lemma 6.2, our problem can be reduced to the case $N = 1$ with $f(y, z) = y^3 + z^5$. To get an explicit basis of the Kuranishi space T^1 type:

$$\begin{aligned} &> \text{F} := y^3 + z^5; \\ &> \text{TyB} := \text{TyurinaBasis(F)}; \\ &\quad \text{TyB} := [yz^3, z^3, yz^2, z^2, yz, z, y, 1] \end{aligned}$$

Therefore

$$T^1 \cong \langle 1, y, z, yz, z^2, yz^2, z^3, yz^3 \rangle_{\mathbb{C}}$$

and given $\Lambda = (\lambda_0, \lambda_1, \dots, \lambda_7) \in T^1$, the associated deformation of U_0 is

$$\begin{aligned} U_{\Lambda} &= \{f_{\Lambda}(y, z) = 0\} \quad \text{where} \quad f_{\Lambda}(y, z) := \\ &= f(y, z) + \lambda_0 + \lambda_1 y + \lambda_2 z + \lambda_3 yz + \lambda_4 z^2 + \lambda_5 yz^2 + \lambda_6 z^3 + \lambda_7 yz^3 \end{aligned}$$

A solution of the jacobian system of partial derivatives is then given by a solution $p_{\Lambda} = (y_{\Lambda}, z_{\Lambda})$ of the following polynomial system in $\mathbb{C}[\lambda][y, z]$

$$(79) \quad \begin{cases} 3y^2 + \lambda_1 + \lambda_3 z + \lambda_5 z^2 + \lambda_7 z^3 = 0 \\ 5z^4 + \lambda_2 + 2\lambda_4 z + 3\lambda_6 z^2 + y(\lambda_3 + 2\lambda_5 z + 3\lambda_7 z^2) = 0 \end{cases}$$

giving 8 critical points for f_Λ . Then p_Λ , one of them, is actually a singular point of U_Λ if and only if

$$(80) \quad p_\Lambda \in U_\Lambda \iff \Lambda \in \mathcal{L} \subset T^1 \quad \text{where}$$

$$(81) \quad \mathcal{L} := \{y_\Lambda^3 + z_\Lambda^5 + \lambda_0 + \lambda_1 y_\Lambda + \lambda_2 z_\Lambda + \lambda_3 y_\Lambda z_\Lambda + \lambda_4 z_\Lambda^2 + \lambda_5 y_\Lambda z_\Lambda^2 + \lambda_6 z_\Lambda^3 + \lambda_7 y_\Lambda z_\Lambda^3 = 0\}.$$

After translating $y \mapsto y + y_\Lambda, z \mapsto z + z_\Lambda$, we get

$$\begin{aligned} f_\Lambda(y + y_\Lambda, z + z_\Lambda) &= f(y, z) + \\ &+ (y_\Lambda^3 + z_\Lambda^5 + \lambda_0 + \lambda_1 y_\Lambda + \lambda_2 z_\Lambda + \lambda_3 y_\Lambda z_\Lambda + \lambda_4 z_\Lambda^2 + \lambda_5 y_\Lambda z_\Lambda^2 + \lambda_6 z_\Lambda^3 + \lambda_7 y_\Lambda z_\Lambda^3) \\ &+ (3y_\Lambda^2 + \lambda_1 + \lambda_3 z_\Lambda + \lambda_5 z_\Lambda^2 + \lambda_7 z_\Lambda^3) y \\ &+ (5z_\Lambda^4 + \lambda_2 + 2\lambda_4 z_\Lambda + 3\lambda_6 z_\Lambda^2 + y_\Lambda(\lambda_3 + 2\lambda_5 z_\Lambda + 3\lambda_7 z_\Lambda^2)) z \\ &+ (\lambda_3 + 2\lambda_5 z_\Lambda + 3\lambda_7 z_\Lambda^2) yz + 3y_\Lambda y^2 + (10z_\Lambda^3 + \lambda_4 + \lambda_5 y_\Lambda + 3\lambda_6 z_\Lambda + 3\lambda_7 y_\Lambda z_\Lambda) z^2 \\ &+ (\lambda_5 + 3\lambda_7 z_\Lambda) yz^2 + (10z_\Lambda^2 + \lambda_6 + \lambda_7 y_\Lambda) z^3 + \lambda_7 yz^3 + 5z_\Lambda z^4 \end{aligned}$$

as can be checked by setting $K := y_\Lambda, L := z_\Lambda$ and typing:

```
> T := nops(TyB):
> F[Lambda] := F+sum(lambda[i]*TyB[T-i], i=0 .. T-1)
FΛ := y3 + z5 + λ0 + λ1y + λ2z + λ3yz + λ4z2 + λ5yz2 + λ6z3 + λ7yz3
> z := Z+L: y := Y+K:
> F[Lambda] := collect(F[Lambda], [Y, Z], 'distributed'):
FΛ := Y3 + 3KY2 + λ7YZ3 + (λ5 + 3λ7L)YZ2 + (3λ7L2 + λ3 + 2λ5L)YZ
+ (λ7L3 + λ1 + λ5L2 + λ3L + 3K2)Y + Z5 + 5LZ4 + (λ6 + λ7K + 10L2)Z3
+ (3λ7KL + λ4 + 3λ6L + 10L3 + λ5K)Z2
+ (λ2 + 3λ7KL2 + 5L4 + 3λ6L2 + λ3K + 2λ5KL + 2λ4L)Z
+ K3 + λ6L3 + λ1K + λ2L + L5 + λ7KL3 + λ0 + λ5KL2 + λ4L2 + λ3KL
```

Define:

$$(82) \quad \begin{aligned} v_0 &= \lambda_3 + 2\lambda_5 z_\Lambda + 3\lambda_7 z_\Lambda^2, & v_1 &= 3y_\Lambda \\ v_2 &= 10z_\Lambda^3 + \lambda_4 + \lambda_5 y_\Lambda + 3\lambda_6 z_\Lambda + 3\lambda_7 y_\Lambda z_\Lambda, & v_3 &= \lambda_5 + 3\lambda_7 z_\Lambda \\ v_4 &= 10z_\Lambda^2 + \lambda_6 + \lambda_7 y_\Lambda, & v_5 &= \lambda_7, & v_6 &= 5z_\Lambda \end{aligned}$$

and $\mathcal{V}_k := \{v_k = 0\}$. Then the proof goes on exactly as in the E_7 case until D_5 singularities: precisely by setting

$$(83) \quad \begin{aligned} \mathcal{W}_2 &:= \{4v_1 v_2 - v_0^2 = 0\} \\ \mathcal{W}_3 &:= \{v_1^3 v_4^2 - v_2(v_1 v_3 + v_2)^2 = 0\} \\ \mathcal{V} &:= \{4v_3^3 + 27v_4^2 = 0\} \end{aligned}$$

we have the inclusions' chain

$$T^1 \supset \mathcal{L} \supset \mathcal{W}_2 \supset \mathcal{W}_2^3 \supset \mathcal{V}_1 \cap \mathcal{W}_2^3 = \mathcal{V}_0^2 \supset \mathcal{V} \cap \mathcal{V}_0^2$$

of subsets parameterizing small deformations whose generic fibre is either smooth or admits a singularity of type A_1, A_2, A_3, D_4 and D_5 , respectively, as can be checked by typing

```
> SD := [Y*Z, Y^2, Z^2, Y*Z^2, Z^3, Y*Z^3, Z^4]:
> s := nops(SD): FL := Y^3+Z^5+sum(v[i]*SD[i+1], i = 0 .. s-1)
```

```

FL := Y^3 + Z^5 + v_0 Y Z + v_1 Y^2 + v_2 Z^2 + v_3 Y Z^2 + v_4 Z^3 + v_5 Y Z^3 + v_6 Z^4
> MGB := MilnorGroebnerBasis(FL, {Y, Z}, plex_min(Z, Y)):
> for i to nops(MGB) do LeadingTerm(MGB[i], plex_min(Z, Y))end do
      v_0, Y
      2 v_1, Y
      -4 v_1^2 v_0^2 + 16 v_1^3 v_2, Z
> F2 := Y^3 + Z^5 + (w[1]*Y + w[2]*Z)^2 + sum(v[j]*SD[j+1], j = 3 .. s-1)
      F2 := Y^3 + Z^5 + (w_1 Y + w_2 Z)^2 + v_3 Y Z^2 + v_4 Z^3 + v_5 Y Z^3 + v_6 Z^4
> print(mu = MilnorNumber(F2, {Y, Z}, plex_min(Z, Y)))
      mu = 2
> MGB2 := MilnorGroebnerBasis(F2, {Y, Z}, plex_min(Z, Y)):
> for i to nops(MGB2) do LeadingTerm(MGB2[i], plex_min(Z, Y))end do
      2 w_2 w_1, Y
      2 w_1^2, Y
      -12 w_1^3 w_2^3 - 12 w_1^5 w_2 v_3 + 12 w_1^6 v_4, Z^2
> w[2] := u*w[1] : v[4] := u*(v[3] + u^2): F2
      Y^3 + Z^5 + (w_1 Y + u w_1 Z)^2 + v_3 Y Z^2 + u (v_3 + u^2) Z^3 + v_5 Y Z^3 + v_6 Z^4
> M1 := Milnor(F2, {Y, Z}, plex_min(Z, Y)): print(mu = M1[4]);
      mu = 3
> MGB3 := M1[1] :
> for i to nops(MGB3) do LeadingTerm(MGB3[i], plex_min(Z, Y))end do
      2 u w_1^2, Y
      -8 w_1^4 v_3^2 + 32 w_1^6 v_6 - 32 w_1^6 u v_5 - 48 w_1^4 u^2 v_3 - 72 w_1^4 u^4, Z^3
      2 w_1^2, Y
> w[1] := 0:
> print(mu = MilnorNumber(F2, {Y, Z}, plex_min(Z, Y)),
> tau = TYURINANumber(F2, {Y, Z}, [plex_min(Z, Y), tdeg(Z, Y)]))
      mu = 4, tau = 4
> unassign('w[1]'):unassign('v[4]'):v[0]:=0:v[1]:=0:v[2]:=0:FL
      Y^3 + Z^5 + v_3 Y Z^2 + v_4 Z^3 + v_5 Y Z^3 + v_6 Z^4
> MGB4 := MilnorGroebnerBasis(FL, {Y, Z}, plex_min(Z, Y)):
> for i to nops(MGB4) do LeadingTerm(MGB4[i], plex_min(Z, Y))end do
      2 v_3, Y Z
      -27 v_4^2 - 4 v_3^3, Z^3
      3, Y^2
> v[3] := -3*a^2 : v[4] := -2*a^3 : FL
      Y^3 + Z^5 - 3 a^2 Y Z^2 - 2 a^3 Z^3 + v_5 Y Z^3 + v_6 Z^4
> print(mu = MilnorNumber(FL, {Y, Z}, plex_min(Z, Y)),
> tau = TyurinaNumber(FL, {Y, Z}, plex_min(Z, Y)))

```

```


$$\mu = 5, \tau = 5$$

> v[4] := 2*a^3 : FL

$$Y^3 + Z^5 - 3a^2YZ^2 + 2a^3Z^3 + v_5YZ^3 + v_6Z^4$$

> print(mu = MilnorNumber(FL,{Y,Z},plex_min(Z,Y)),
> tau = TyurinaNumber(FL,{Y,Z},plex_min(Z,Y)))

$$\mu = 5, \tau = 5$$


```

Let us now go on by considering:

```

> MGB5 := MilnorGroebnerBasis(FL, {Y,Z},plex_min(Z,Y)):
> for i to nops(MGB5) do LeadingTerm(MGB5[i],plex_min(Z,Y))end do

$$3, Y^2$$


$$-48a^4v_5 + 48a^3v_6, Z^4$$


$$-6a^2, YZ$$

> a := 0 : FL

$$Y^3 + Z^5 + v_5YZ^3 + v_6Z^4$$

> print(mu = MilnorNumber(FL,{Y,Z},plex_min(Z,Y)),
> tau = TyurinaNumber(FL,{Y,Z},plex_min(Z,Y)))

$$\mu = 6, \tau = 6$$

> MGB6 := MilnorGroebnerBasis(FL,{Y,Z},plex_min(Z,Y)):
> for i to nops(MGB6) do LeadingTerm(MGB6[i],plex_min(Z, Y))end do

$$3, Y^2$$


$$3v_5, YZ^2$$


$$16v_6^2, Z^4$$

> v[5] := 0 : FL

$$Y^3 + Z^5 + v_6Z^4$$

> print(mu = MilnorNumber(FL,{Y,Z},plex_min(Z,Y)),
> tau = TyurinaNumber(FL,{Y,Z},plex_min(Z, Y)))

$$\mu = 6, \tau = 6$$

> unassign('v[5]') : v[6] := 0 : FL

$$Y^3 + Z^5 + v_5YZ^3$$

> print(mu = MilnorNumber(FL,{Y,Z},plex_min(Z,Y)),
> tau = TyurinaNumber(FL,{Y,Z},plex_min(Z,Y)))

$$\mu = 7, \tau = 7$$


```

meaning that

- $0 \in f_{\Lambda}^{-1}(0)$ is a singularity of type E_6 for Λ generic in $\mathcal{V}_0^4$,
- $0 \in f_{\Lambda}^{-1}(0)$ is a singularity of type E_7 for Λ generic in $\mathcal{V}_0^4 \cap \mathcal{V}_6$.

On the other hand, by considering the second leading coefficient in MGB5, define the following codimension 2 subset of $\mathcal{L}$

$$(84) \quad \mathcal{V}' := \{v_3v_5^2 + 3v_6^2 = v_4v_5^3 + 2v_6^3 = 0\} \subset \mathcal{V}$$

and type

```

> unassign('v[6]') : unassign('a') : FL

```

```


$$Y^3 + Z^5 - 3a^2YZ^2 - 2a^3Z^3 + v_5YZ^3 + v_6Z^4$$

> v[6] := a*v[5] : FL

$$Y^3 + Z^5 - 3a^2YZ^2 - 2a^3Z^3 + v_5YZ^3 + av_5Z^4$$

> print(mu = MilnorNumber(FL,{Y,Z},plex_min(Z,Y)),
> tau = TyurinaNumber(FL,{Y,Z},plex_min(Z,Y)))

$$\mu = 6, \tau = 6$$

> MGB6b := MilnorGroebnerBasis(FL,{Y,Z},plex_min(Z,Y)):
> for i to nops(MGB6b) do LeadingTerm(MGB6b[i],plex_min(Z,Y))end do

$$3, Y^2$$


$$-6a^2, YZ$$


$$60a^3v_5 + 5a^2v_5^3, Z^5$$

> a:=0 : FL

$$Y^3 + Z^5 + v_5YZ^3$$

> unassign('a') : a:=(1/12)*v[5]^2 : FL

$$Y^3 + Z^5 - 1/48v_5^4YZ^2 + \frac{1}{864}v_5^6Z^3 + v_5YZ^3 - 1/12v_5^3Z^4$$

> print(mu = MilnorNumber(FL,{Y,Z},plex_min(Z,Y)),
> tau = TyurinaNumber(FL,{Y,Z},plex_min(Z,Y)))

$$\mu = 7, \tau = 7$$


```

Consider the latter leading coefficient in MGB6b and define ¹⁰ the following codimension 3 subset of $\mathcal{L}$

$$(85) \quad \mathcal{V}'' := \mathcal{V}' \cap \{12v_6 + v_5^3 = 0\} ,$$

Then, by Theorem 6.3(3), we get that

- $0 \in f_{\Lambda}^{-1}(0)$ is a singularity of type D_6 for Λ generic in $\mathcal{V}' \cap \mathcal{V}_0^2$,
- $0 \in f_{\Lambda}^{-1}(0)$ is a singularity of type D_7 for Λ generic in $\mathcal{V}'' \cap \mathcal{V}_0^2$,

while $\mathcal{V}' \cap \mathcal{V}_0^3 = \mathcal{V}_0^4 \cap \mathcal{V}_6$ and we get the E_7 singularities already discussed. A further specialization here gives the trivial deformation.

Let us then come back to consider $\mathcal{W}_2^3$ and look at the second leading coefficient in MGB3. Define

$$(86) \quad \mathcal{W}_4 := \{16v_1^5v_2v_5^2 - [(v_1v_3 + 3v_2)^2 - 4v_1^3v_6]^2 = 0\}$$

whose equation is obtained by eliminating u and t from the following set of equations, parameterizing $\mathcal{W}_3^4 \setminus \mathcal{V}_1^2$,

$$(87) \quad v_2 - u^2v_1 = 0, \quad v_4 - u(v_3 + u^2) = 0, \quad 4v_1(v_6 - uv_5) - (v_3 + 3u^2)^2 = 0.$$

Observe that $\mathcal{V}_1^2 \subseteq \mathcal{W}_3^4 \Rightarrow \mathcal{V}_0^2 \subseteq \mathcal{W}_2^4$ and define the following codimension 1 Zariski closed subset ¹¹ of $\mathcal{W}_2^3$

$$(88) \quad \widetilde{\mathcal{W}}_2^4 := \overline{\mathcal{W}_2^4 \setminus \mathcal{V}_0^2}.$$

Then ¹²

¹⁰The reader may check that choosing $v_4 = 2a^3$, $v_6 = -av_5$, $12a = v_5^2$ leads to same results on Milnor and Tyurina numbers and to the same condition defining $\mathcal{V}''$ in (85).

¹¹The same considerations explained by footnote 6 are still holding, here.

¹²The reader may check that the choice $v_3 = -2tw_1 - 3u^2$ leads to the same conclusion.

```

> v[6]:=t^2+u*v[5] : v[3]:=2*t*w[1]-3*u^2 : F2
      Y^3 + Z^5 + (w_1 Y + u w_1 Z)^2 + (2 t w_1 - 3 u^2) Y Z^2 +
      u (2 t w_1 - 2 u^2) Z^3 + v_5 Y Z^3 + (t^2 + u v_5) Z^4
> print(mu = MilnorNumber(F2,{Y,Z},plex_min(Z,Y)),
> tau = TYURINANumber(F2,{Y,Z},[plex_min(Z,Y), tdeg(Z,Y)]))
      μ = 4, τ = 4

```

and Theorem 6.3(2) allows to conclude that

- $0 \in f_{\Lambda}^{-1}(0)$ is a singularity of type A_4 for Λ generic in $\widetilde{\mathcal{W}}_2^4$.

Then type:

```

> MGB4b := MilnorGroebnerBasis(F2,{Y,Z},plex_min(Z,Y)):
> for i to nops(MGB4b) do LeadingTerm(MGB4b[i],plex_min(Z,Y))end do
      120 t^3 w_1^4 u + 40 t^2 w_1^5 v_5 - 40 t w_1^6, Z^4
      2 u w_1^2, Y
      2 w_1^2, Y
> w[1]:=0 : F2
      Y^3 + Z^5 - 3 u^2 Y Z^2 - 2 u^3 Z^3 + v_5 Y Z^3 + (t^2 + u v_5) Z^4
> print(mu = MilnorNumber(F2,{Y,Z},plex_min(Z,Y)),
> tau = TyurinaNumber(F2,{Y,Z},plex_min(Z,Y)))
      μ = 5, τ = 5

```

This means that $\mathcal{V}_1 \cap \widetilde{\mathcal{W}}_2^4 = \mathcal{V} \cap \mathcal{V}_0^2$ obtaining a generic D_5 singularity, as already described above. Go on:

```

> unassign('w[1]') : t:=0 : F2
      Y^3 + Z^5 + (w_1 Y + u w_1 Z)^2 - 3 u^2 Y Z^2 - 2 u^3 Z^3 + v_5 Y Z^3 + u v_5 Z^4
> print(mu = MilnorNumber(F2,{Y,Z},plex_min(Z,Y)),
> tau = TYURINANumber(F2,{Y,Z},[plex_min(Z,Y), tdeg(Z,Y)]))
      μ = 4, τ = 4
> unassign('t') : u:=0 : F2
> print(mu = MilnorNumber(F2,{Y,Z},plex_min(Z,Y)),
> tau = TYURINANumber(F2,{Y,Z},[plex_min(Z,Y), tdeg(Z,Y)]))
      μ = 4, τ = 4

```

Then we have to consider the relation assigned by the first leading coefficient in MGB4b, giving the further condition

$$v_1 - w_1 v_5 t - 3 u t^2 = 0 ,$$

which added to (87) parameterizes the following (reducible) codimension 4 Zariski closed subset of $\mathcal{L}$

$$(89) \quad \widetilde{\mathcal{W}}_2^5 := \overline{\mathcal{W}_2^5 \setminus \mathcal{V}_0^2}$$

where ¹³

$$\mathcal{W}_5 := \{[4v_1^5v_5^2(v_1v_3 + 3v_2)^2 + 16v_1^7(v_1^2 + v_1v_3v_5 + 3v_2v_5) - 9v_2(v_1v_3 + 3v_2)^4] \cdot$$

$$(90) \quad [4v_1^5v_5^2(v_1v_3 + 3v_2)^2 + 16v_1^7(v_1^2 - v_1v_3v_5 - 3v_2v_5) - 9v_2(v_1v_3 + 3v_2)^4] = 0\}$$

Since

$$\begin{aligned} &> \text{u} := 1/12 * (\text{b}^2 - \text{v}[5]^2) : \quad \text{w}[1] := (1/2) * \text{t} * (\text{v}[5] - \text{b}) : \quad \text{F2} \\ &Y^3 + Z^5 + (1/2 t (v_5 - b) Y + 1/2 (1/12 b^2 - 1/12 v_5^2) t (v_5 - b) Z)^2 + \\ &\quad \left(t^2 (v_5 - b) - 3 (1/12 b^2 - 1/12 v_5^2)^2 \right) Y Z^2 + \\ &\quad (1/12 b^2 - 1/12 v_5^2) \left(t^2 (v_5 - b) - 2 (1/12 b^2 - 1/12 v_5^2)^2 \right) Z^3 + v_5 Y Z^3 + \\ &\quad (t^2 + (1/12 b^2 - 1/12 v_5^2) v_5) Z^4 \\ &> \text{M5} := \text{Milnor}(\text{F2}, \{Y, Z\}, \text{plex_min}(Z, Y)) : \text{print}(\mu = \text{M5}[4]) \\ &\quad \mu = 5 \end{aligned}$$

then Theorem 6.3 (2) gives that ¹⁴

- $0 \in f_{\Lambda}^{-1}(0)$ is a singularity of type A_5 for Λ generic in $\widetilde{\mathcal{W}}_2^5$.

Furthermore:

$$\begin{aligned} &> \text{MGB5} := \text{M5}[1] : \text{for } i \text{ to } \text{nops}(\text{MGB5}) \text{ do} \\ &> \text{factor}(\text{LeadingTerm}(\text{MGB5}[i], \text{plex_min}(Z, Y))) \text{end do} \\ &\quad \frac{1594323}{576460752303423488} t^6 (b - v_5)^3 (b^3 - b^2 v_5 - 8 t^2), Z^5 \\ &\quad 1/2 t^2 (v_5 - b)^2 (1/12 b^2 - 1/12 v_5^2), Y \\ &\quad 1/2 t^2 (v_5 - b)^2, Y \\ &> \text{t} := 0 : \quad \text{F2} \\ &Y^3 + Z^5 - 3 (1/12 b^2 - 1/12 v_5^2)^2 Y Z^2 - 2 (1/12 b^2 - 1/12 v_5^2)^3 Z^3 + \\ &\quad v_5 Y Z^3 + (1/12 b^2 - 1/12 v_5^2) v_5 Z^4 \\ &> \text{M5b} := \text{Milnor}(\text{F2}, \{Y, Z\}, \text{plex_min}(Z, Y)) : \text{print}(\mu = \text{M5b}[4]) \\ &\quad \mu = 6 \\ &> \text{T5b} := \text{TYURINA}(\text{F2}, \{Y, Z\}, [\text{plex_min}(Z, Y), \text{tdeg}(Z, Y)]) : \text{print}(\tau = \text{T5b}[4]) \\ &\quad \tau = 6 \end{aligned}$$

Observe that, by identifying $a = (b^2 - v_5^2)/12$, we are precisely dealing with deformations parameterized by the above considered algebraic closed subset $\mathcal{V}' \cap \mathcal{V}_0^2$ of $\mathcal{L}$, whose generic fibre has a singularity of type D_6 . On the other hand

$$\begin{aligned} &> \text{unassign}('t') : \quad \text{v}[5] := -b : \quad \text{F2} \\ &\quad Y^3 + Z^5 + t^2 b^2 Y^2 - 2 t^2 b Y Z^2 - b Y Z^3 + t^2 Z^4 \\ &> \text{M5c} := \text{Milnor}(\text{F2}, Y, Z, \text{plex_min}(Z, Y)) : \text{print}(\mu = \text{M5c}[4]) \\ &\quad \mu = 5 \\ &> \text{v}[5] := b : \quad \text{F2} \\ &\quad Y^3 + Z^5 + b Y Z^3 + t^2 Z^4 \end{aligned}$$

¹³The following equation of $\mathcal{W}_5$ is obtained by carefully applying Maple's commands `eliminate` and `EliminationIdeal` in the `PolynomialIdeals` package.

¹⁴The reader may check that setting $w_1 := t(v_5 + b)/2$ leads to the same conclusion.

still giving the above considered deformations parameterized by $\mathcal{V}_0^4$ whose generic fibre has a singularity of type E_6 . Let us then consider the relation given by the last factor in the first leading coefficient of MGB5:

```

> t:=c*b : v[5]:=b-8*c^2 : F2

$$Y^3 + Z^5 + \left(-4c^3bY - 4\left(1/12b^2 - 1/12(b-8c^2)^2\right)c^3bZ\right)^2 +$$


$$\left(-8c^4b^2 - 3\left(1/12b^2 - 1/12(b-8c^2)^2\right)^2\right)YZ^2 +$$


$$\left(1/12b^2 - 1/12(b-8c^2)^2\right)\left(-8c^4b^2 - 2\left(1/12b^2 - 1/12(b-8c^2)^2\right)^2\right)Z^3 +$$


$$(b-8c^2)YZ^3 + \left(c^2b^2 + \left(1/12b^2 - 1/12(b-8c^2)^2\right)(b-8c^2)\right)Z^4$$

> M5d := Milnor(F2, {Y,Z}, plex_min(Z,Y)) : print(mu = M5d[4])

$$\mu = 6$$

> T5d := TYURINA(F2, {Y,Z}, [plex_min(Z,Y), tdeg(Z,Y)]) :
> print(tau = T5d[4])

$$\tau = 6$$

> MGB6a := M5d[1] : for i to nops(MGB6a) do
> LeadingTerm(MGB6a[i], plex_min(Z, Y)) end do:
> for i to nops(MGB6a) do
> factor(LeadingCoefficient(MGB6a[i], plex_min(Z, Y))) end do

$$32c^6b^2$$


$$-3670016c^{16}b^8(b+8c^2)(b+16c^2)^2$$


$$\frac{128}{3}c^8b^2(-4c^2+b)$$

> c := 0 : F2

$$Y^3 + Z^5 + bYZ^3$$

> unassign('c') : b:=0 : F2

$$Y^3 + Z^5 - \frac{256}{3}YZ^2c^8 + \frac{8192}{27}Z^3c^{12} - 8YZ^3c^2 + \frac{128}{3}Z^4c^6$$

> M6a := Milnor(F2, {Y,Z}, plex_min(Z,Y)) : print(mu = M6a[4])

$$\mu = 7$$

> T6a := TYURINA(F2, {Y,Z}, [plex_min(Z,Y), tdeg(Z,Y)]) :
> print(tau = T6a[4]) :

$$\tau = 7$$

> b := 4*c^2 : F2

$$Y^3 + Z^5 + 256Y^2c^{10} - 128YZ^2c^8 - 4YZ^3c^2 + 16Z^4c^6$$

> M6b := Milnor(F2, {Y,Z}, plex_min(Z, Y)) : print(mu = M6b[4])

$$\mu = 6$$

> T6b := TYURINA(F2, {Y,Z}, [plex_min(Z, Y), tdeg(Z, Y)]) :
> print(tau = T6b[4])

$$\tau = 6$$

> b := -8*c^2 : F2

$$Y^3 + Z^5 + (32c^5Y - 512c^9Z)^2 - 1280YZ^2c^8 + 16384Z^3c^{12} - 16YZ^3c^2 + 320Z^4c^6$$

> M6c := Milnor(F2, {Y,Z}, plex_min(Z, Y)) : print(mu = M6c[4])

```

```

                                 $\mu = 7$ 
> T6c := TYURINA(F2, {Y,Z}, [plex_min(Z, Y), tdeg(Z, Y)]) : print(tau
= T6c[4])
                                 $\tau = 7$ 
> b := -16*c^2 : F2
                                
$$Y^3 + Z^5 + \left(64c^5Y - \frac{5120}{3}c^9Z\right)^2 - \frac{12544}{3}YZ^2c^8 +$$

                                
$$\frac{2498560}{27}Z^3c^{12} - 24YZ^3c^2 + 896Z^4c^6$$

> M6d := Milnor(F2, {Y,Z}, plex_min(Z, Y)) : print(mu = M6d[4])
                                 $\mu = 6$ 
> T6d := TYURINA(F2, {Y,Z}, [plex_min(Z, Y), tdeg(Z, Y)]) :
> print(tau = T6d[4])
                                 $\tau = 6$ 

```

This means that:

- equation $b^3 - b^2v_5 - 8t^2 = 0$ defines the following codimension 5 (reducible) algebraic subset of $\mathcal{L}$

$$(91) \quad \widetilde{\mathcal{W}}_2^6 := \mathcal{W}_6 \cap \widetilde{\mathcal{W}}_2^5 ,$$

where ¹⁵

$$(92) \quad \begin{aligned} \mathcal{W}_6 := & \{ [32v_1^9 - 2v_1^5v_5(v_1v_3 + 3v_2)(8v_1^2 - 3v_2v_5 - v_1v_3v_5) + (v_1v_3 + 3v_2)^5] \cdot \\ & [32v_1^9 + 2v_1^5v_5(v_1v_3 + 3v_2)(8v_1^2 + 3v_2v_5 + v_1v_3v_5) - (v_1v_3 + 3v_2)^5] = 0 \} \end{aligned}$$

whose generic point Λ is such that $0 \in f_{\Lambda}^{-1}(0)$ is an A_6 singularity,

- setting $c = 0$ means studying deformations parameterized by $\mathcal{V}_0^4 \cap \mathcal{V}_6$, generically admitting an E_7 singularity, as already observed above,
- and setting $b = 0$ means studying deformations parameterized by $\mathcal{V}_0^2 \cap \mathcal{V}''$, generically admitting a D_7 singularity and already considered above, too,
- the further step is then to consider the relation $b + 8c^2 = 0$ defining the following codimension 6 (reducible) algebraic subset of $\mathcal{L}$

$$(93) \quad \widetilde{\mathcal{W}}_2^7 := \mathcal{W}_7 \cap \widetilde{\mathcal{W}}_2^6 \quad \text{with} \quad \mathcal{W}_7 := \{256v_2 - v_1v_5^4 = 0\}$$

whose generic point Λ is such that $0 \in f_{\Lambda}^{-1}(0)$ is an A_7 singularity.

The check of relations (78) are then left to the reader. $\square$

6.7. A list of very special adjacencies. As a consequence of the analysis performed in the previous sections, we are now able to concretely write down some very special small 1-parameter deformations of a A_n , D_n , E_6 , E_7 or E_8 , realizing adjacencies not directly mentioned in [3] and [5] (except for those in 6.7.1). The 1-parameter deformations we are going to list in the following are obtained by last steps in proofs of Theorems 6.5, 6.6, 6.8, 6.10 and 6.11, giving precisely 1-parameter deformations, after some possible parameter's re-scaling.

¹⁵The following equation of $\mathcal{W}_6$ is obtained by carefully applying Maple's commands `eliminate` and `EliminationIdeal` in the `PolynomialIdeals` package.

6.7.1. $A_{n-1} \longleftarrow D_n$. Assume that $n \geq 4$ is either $n = 2m + 4$ when n is even, or $n = 2m + 5$ when odd. Then consider the 1-parameter family $\mathcal{X}_t := \{f_t(\mathbf{x}) = 0\}$, $t \in \mathbb{C}$, where

$$f_0(\mathbf{x}) := \sum_{i=1}^{N-1} x_i^2 + x_N^2 x_{N+1} + x_{N+1}^{n-1}$$

and either

$$f_t(\mathbf{x}) := f_0(\mathbf{x}) + t(x_N + it^m x_{N+1})^2 + \sum_{k=3}^{2m+2} (-t)^{2m+3-k} x_{N+1}^k \quad (n \text{ even})$$

or

$$f_t(\mathbf{x}) := f_0(\mathbf{x}) + t^2(x_N + t^{2m+1} x_{N+1})^2 + \sum_{k=3}^{2m+3} (-t^2)^{2m+4-k} x_{N+1}^k \quad (n \text{ odd}).$$

Then $\mathcal{X}_0$ is an isolated D_n point and, for generic $t \in \mathbb{C}$, $\mathcal{X}_t$ admits the unique singular point $\mathbf{0} \in f_t^{-1}(0)$ which is of type A_{n-1} .

6.7.2. $A_5 \longleftarrow E_6$. Consider the 1-parameter family $\mathcal{X}_t := \{f_t(\mathbf{x}) = 0\}$, $t \in \mathbb{C}$, with

$$\begin{aligned} f_0(\mathbf{x}) &:= \sum_{i=1}^{N-1} x_i^2 + x_N^3 + x_{N+1}^4 \\ f_t(\mathbf{x}) &:= f_0(\mathbf{x}) + t^2 x_N^2 + 2t x_N x_{N+1}^2. \end{aligned}$$

Then $\mathcal{X}_0$ is an isolated E_6 singular point and, for generic $t \in \mathbb{C}$, $\mathcal{X}_t$ admits the unique singular point $\mathbf{0} \in f_t^{-1}(0)$ which is of type A_5 .

6.7.3. $D_5 \longleftarrow E_6$. Let $f_0(\mathbf{x})$ be as in the previous case and assume

$$f_t(\mathbf{x}) := f_0(\mathbf{x}) - 3t^2 x_N x_{N+1}^2 - 2t^3 x_{N+1}^3.$$

Then, for generic $t \in \mathbb{C}$, $\mathcal{X}_t$ admits the unique singular point $\mathbf{0} \in f_t^{-1}(0)$ which is of type D_5 .

6.7.4. $A_6 \longleftarrow E_7$. Consider the 1-parameter family $\mathcal{X}_t := \{f_t(\mathbf{x}) = 0\}$, $t \in \mathbb{C}$, with

$$f_0(\mathbf{x}) := \sum_{i=1}^{N-1} x_i^2 + x_N^3 + x_N x_{N+1}^3$$

$$f_t(\mathbf{x}) := f_0(\mathbf{x}) + 432t^3(x_N + 4tx_{N+1})^2 - 120t^2 x_N x_{N+1}^2 - 416t^3 x_{N+1}^3 + 7tx_{N+1}^4$$

Then $\mathcal{X}_0$ is an isolated E_7 singular point and, for generic $t \in \mathbb{C}$, $\mathcal{X}_t$ admits the unique singular point $\mathbf{0} \in f_t^{-1}(0)$ which is of type A_6 .

6.7.5. $D_6 \longleftarrow E_7$. Let $f_0(\mathbf{x})$ be as in the previous case and assume

$$f_t(\mathbf{x}) := f_0(\mathbf{x}) - 3t^2 x_N x_{N+1}^2 - 2t^3 x_{N+1}^3 + tx_{N+1}^4.$$

Then, for generic $t \in \mathbb{C}$, $\mathcal{X}_t$ admits the unique singular point $\mathbf{0} \in f_t^{-1}(0)$ which is of type D_6 .

6.7.6. $A_7 \longleftarrow E_8$. Consider the 1-parameter family $\mathcal{X}_t := \{f_t(\mathbf{x}) = 0\}$, $t \in \mathbb{C}$, with

$$f_0(\mathbf{x}) := \sum_{i=1}^{N-1} x_i^2 + x_N^3 + x_{N+1}^5$$

$$f_t(\mathbf{x}) := f_0(\mathbf{x}) + t^5(x_N - t^2 x_{N+1})^2 - 5t^4 x_N x_{N+1}^2 + 4t^6 x_{N+1}^3 - 4t x_N x_{N+1}^3 + 5t^3 x_{N+1}^4$$

Then $\mathcal{X}_0$ is an isolated E_8 singular point and, for generic $t \in \mathbb{C}$, $\mathcal{X}_t$ admits the unique singular point $\mathbf{0} \in f_t^{-1}(0)$ which is of type A_7 .

6.7.7. $D_7 \longleftarrow E_8$. Let $f_0(\mathbf{x})$ be as in the previous case and assume

$$f_t(\mathbf{x}) := f_0(\mathbf{x}) - 27t^4 x_N x_{N+1}^2 + 54t^6 x_{N+1}^3 - 6t x_N x_{N+1}^3 + 18t^3 x_{N+1}^4.$$

Then, for generic $t \in \mathbb{C}$, $\mathcal{X}_t$ admits the unique singular point $\mathbf{0} \in f_t^{-1}(0)$ which is of type D_7 .

REFERENCES

- [1] Soon available at <http://www.maplesoft.com/applications>
- [2] <http://www.singular.uni-kl.de>
- [3] Arnol'd V. I. "Local normal forms of functions" *Invent. Math.* **35** (1976), 87–109.
- [4] Arnol'd V. I., Goryunov V. V., Lyashko O. V. and Vasil'ev V. A. *Singularity theory I*, translated from the 1988 Russian original by A. Iacob, reprint of the original English edition from the series Encyclopaedia of Mathematical Sciences [Dynamical systems. VI, Encyclopaedia Math. Sci., 6, Springer, Berlin, 1993; MR1230637 (94b:58018)] Springer-Verlag, Berlin, 1998.
- [5] Arnol'd V. I., Gusein-Zade S. M. and Varchenko A. N. *Singularities of differentiable maps*, Vol. I, *The classification of critical points caustics and wave fronts*, Monographs in Mathematics **82**, Birkhäuser Boston, Inc., Boston, MA, 1988.
- [6] Barth W., Peters C. and Van de Ven A. *Compact complex surfaces* vol. **4** E.M.G., Springer-Verlag (1984)
- [7] Cox D., Little J. and O'Shea D. *Ideals, varieties, and algorithms. An introduction to computational algebraic geometry and commutative algebra*. Third edition, Undergraduate Texts in Mathematics, Springer, New York (2007).
- [8] Decker W. and Lossen C. *Computing in algebraic geometry. A quick start using SINGULAR* Algorithms and Computation in Mathematics **16**, Springer-Verlag, Berlin; Hindustan Book Agency, New Delhi (2006).
- [9] Douady A. "Le problème des modules locaux pour les espaces C-analytiques compacts" *Ann. scient. Éc. Norm. Sup.* 4e série, **7** 569–602 (1974).
- [10] Godement R. *Topologie Algébrique et Théorie des Faisceaux*, Hermann, Paris (1958).
- [11] Grauert H. "Der Satz von Kuranishi für Kompakte Komplexe Räume" *Invent. Math.* **25**, 107–142 (1974).
- [12] Greuel G.-M., Lossen C. and Shustin E. *Introduction to singularities and deformations* Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2007.
- [13] Greuel G.-M. and Pfister G. *A Singular introduction to commutative algebra* Springer-Verlag, Berlin, 2002.
- [14] Grothendieck A. "Sur quelques points d'algèbre homologique" *Tôhoku Math. J.* **9**, 199–221 (1957).
- [15] Lichtenbaum S. and Schlessinger M. "On the cotangent complex of a morphism" *Trans. A.M.S.* **128**, 41–70 (1967).
- [16] E. J. N. Looijenga *Isolated singular points on complete intersections* London Mathematical Society Lecture Note Series, **77**, Cambridge University Press, Cambridge, 1984.
- [17] E. J. N. Looijenga and J. Steenbrink "Milnor number and Tjurina number of complete intersections" *Math. Ann.* **271** (1) (1985), 121–124.
- [18] Milnor J. *Singular points of complex hypersurfaces*, Annals of Math. Studies **61**, Princeton University Press, Princeton (1968).
- [19] Milnor J. and Orlik P. "Isolated singularities defined by weighted homogeneous polynomials", *Topology* **9** (1970), 385–393.

- [20] Mora F. “An algorithm to compute the equations of tangent cones” in *Computer algebra (Marseille, 1982)*, Lecture Notes in Comput. Sci. **144**, Springer Berlin-New York (1982), 158–165.
- [21] Palamodov V. P. “The existence of versal deformations of complex spaces” *Dokl. Akad. Nauk SSSR* **206** (1972), 538–541.
- [22] Palamodov V. P. “Deformations of complex spaces” *Russian Math. Surveys* **31**(3) (1976), 129–197; from russian *Uspekhi Mat. Nauk* **31**(3) (1976), 129–194.
- [23] Reid M. “Canonical 3-folds” in *Journées de géométrie algébrique d’Angers*, Sijthoff & Nordhoff (1980), 671–689.
- [24] Saito K. “Quasihomogene isolierte Singularitäten von Hyperflächen” *Invent. Math.* **14** (1971), 123–142.
- [25] J. Stevens *Deformations of singularities* Lecture Notes in Mathematics **1811**, Springer-Verlag, Berlin, 2003.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI TORINO, VIA CARLO ALBERTO 10, 10123 TORINO

E-mail address: `michele.rossi@unito.it`, `lea.terracini@unito.it`